

# Adiposity, joint and systemic inflammation: the additional risk of having a metabolic syndrome in rheumatoid arthritis

Gianfranco Ferraccioli, Elisa Gremese

Division of Rheumatology, Catholic University of the Sacred Heart, Rome, Italy

## Summary

Adiposity is a predisposing condition to atherosclerosis, and rheumatoid arthritis (RA) also predisposes to accelerated atherosclerosis. Adiposity is one of the key features of the metabolic syndrome (MetS) and it is well recognised that a metabolic syndrome (and fat tissue) is a major player in this complex network. Endothelial dysfunction and carotid intima-media thickness, early pre-clinical markers of atherosclerosis which are the main determinants of cardiovascular (CV) morbidity and mortality, occur early on in RA. RA patients have an incidence of CV diseases at least two times higher than the general population.

MetS and RA have a low and a severe-moderate degree of inflammation in common, respectively. Adipose tissue has emerged as a dynamic organ that releases several inflammatory and immune mediators (adipokines). In addition, fat has been recognised as a producer of B cell activating factor (BAFF) and of chemerin, an inducer at the chondrocyte level of IL1 $\beta$ , TNF $\alpha$ , IL6, IL8 and MMP13, thus possibly contributing to cartilage damage. Since fat produces inflammation, to obtain a full control of the CV risk in RA, data suggest that it is therefore mandatory to have a “tight control” of both RA and MetS-related inflammation, especially if RA presents MetS as a co-morbidity.

**Key words:** *rheumatoid arthritis; adiposity; adipokines; chemerin; remission*

## Introduction

Rheumatoid arthritis (RA) pathobiology seems to share some common pathways with atherosclerosis, including endothelial dysfunction which is related to underlying chronic inflammation and presents in the early phases of the disease [1–4]. A possible aggravating risk is represented by the co-existence of a metabolic syndrome (MetS), characterised by a combination of various risk factors that imply additional cardiovascular morbidity that is greater than the sum of the risks associated with each individual component. Several studies have demonstrated that inflammatory processes are involved in the pathogenesis of the metabolic syndrome. On the other hand, there is evidence

that components of cardiovascular (CV) risk increase the inflammatory burden in RA.

## Cardiovascular risk in RA

A recent, Dutch, cross-sectional study found that age- and gender-adjusted odds ratios for CV diseases derived from these cohorts were 3.1 for RA patients and 2.3 for individuals with type 2 diabetes (T2DM) with respect to healthy subjects, indicating that the CV risk in RA is comparable to diabetes, one of the most relevant CV risk factors. It also showed a prevalence of CV events (coronary, cerebral and peripheral arterial diseases) of 13% in non diabetic RA patients, 12% in subjects with T2DM and 5% in non diabetic individuals [5].

The expected life span of patients with RA is known to be shorter than in healthy controls [6], with a standardised overall mortality ratio of between 1.3 and 3 compared to the general population [7], and it is well recognised that, as in the general population, cardiovascular diseases are the leading cause of death. Large epidemiological studies from the last several decades have confirmed that patients with RA are 30 to 60% more likely to suffer a CV event than subjects from the general population [1, 7].

To assess the incidence of CV events, the Dutch authors prospectively followed two cohorts of 263 non diabetic RA patients and 1492 non diabetic individuals for 3 years [8, 9] and they confirmed that patients with RA have about a 2-fold higher CV diseases risk than the general population. The magnitude of the increased CV risk occurred was at least as that of T2DM patients, and this confirmed the previous data in the literature [10].

Histological data provided evidence that coronary arteries from autopsied RA patients have more inflammation in the media and adventitia and more fragile atherosclerotic plaques, but less atherosclerosis, when compared to coronary arteries from age- and sex-matched controls who died from CV diseases. Despite this, the number of acute coronary lesions and grades of stenosis were similar [11, 12, 13]. A possible interpretation of these differences is that the mechanisms responsible for cardiovascular morbidity and mortality are likely to be different in patients with RA.

When comparing the dramatic improvements in the overall mortality rates of the general U.S. population (from 1.0–1.2 to 0.2–0.3), it seems that RA patients have not experienced improvements in survival over the past 4 decades (1965–2005 mortality rates: 2.4–2.5) despite the progress made in the diagnosis and therapy of RA. Clear-cut data suggest that there is a widening mortality gap between RA patients and the general population [14]. In agreement with this analysis, a retrospective population-based cohort study suggested that, even at the moment of RA diagnosis, the absolute CV risk in RA patients was similar to that of subjects without RA who were 5–10 years older [15].

These data suggest that mortality has not really been modified during the past 50 years [16]. Of note, we must take into account that most of these studies considered patients with a diagnosis made before the 1990s, when disease-modifying anti-rheumatic drugs (DMARDs) were poorly prescribed and only minimal attention was paid to the control of inflammation and the prevention of cardiovascular events. It will certainly be interesting to assess the effects of an early diagnosis, of a “tight control” of disease activity and of the new therapies, in particular of the biological drugs, on CV risk. In agreement with the hypothesis that control of CV risk relies on better disease activity control, preliminary evidence reports an improvement of overall cause mortality, and also of CV mortality, when patients were treated very early on [17] and with a more aggressive therapy [18, 19]. The conclusions are still controversial and more data are really needed, especially because a very recent report showed that the risk of cardiovascular events and survival in patients who received TNF- $\alpha$  antagonists was not different from those who received other DMARDs (with a CV morbidity of 38% in both groups) [20]. The strength of this work was that it was conducted in a cohort of about 20'800 U.S. veterans who were diagnosed with RA over a period of seven years, although the bias is that the data were collected retrospectively from an administrative registry. Data from further prospective long-term studies and in early arthritis registries could really help to elucidate this fundamental issue.

### Metabolic syndrome and RA

The prevalence of Metabolic syndrome (MetS) has been assessed in patients with many systemic rheumatic diseases [21]. Recently, Karvounaris *et al* found a prevalence of MetS (defined with NCEP ATP III criteria) in RA patients (40%), comparable with their control population, and documented a relationship between disease activity and the presence of MetS [22]. Another study showed that MetS was significantly more prevalent in American patients with long-standing RA (42% – WHO and NCEP/ATPIII criteria) as well as in early RA patients (31% and 30% – WHO and NCEPIII criteria respectively) than in controls (11% and 22% – WHO and NCEP/ATPIII criteria, respectively) [23].

The same problem can be seen from the alternative angle, and indeed Karvounaris *et al* found that the risk of having a moderate-to-severe RA was higher in patients with metabolic syndrome than in those without, and that disease activity correlated with the number of MetS parameters

present [22]. The take-home message might be that MetS has an inflammatory milieu leading to the occurrence of a more severe RA. If this were true, the therapeutic response should differ in RA carriers of MetS versus non carriers.

Up to now no data are available in the literature on the possible effect of MetS on the response to therapy. Our group evaluated a cohort of 115 long standing RA patients with an active disease, for the presence of metabolic syndrome and disease activity at the start of an anti-TNF $\alpha$  drug prescription and after 12 months of follow-up (tables 1 and 2). At baseline, MetS was seen in 27% of our patients. Over time there was a statistically significant increase in the levels of total cholesterol (mean value 184.1 mg/dl at T0 vs. 192 mg/dl at T12) and triglycerides (mean value 122.3 mg/dl at T0 vs. 134.5 mg/dl at T12) without changes of glucose, high-density lipoprotein (HDL), artery pressure levels or body mass index (BMI). Importantly, we observed that RA patients with a metabolic syndrome at baseline had a significantly lower chance to achieve a good response (DAS<2.4) at 1 year of follow-up, compared to patients without a MetS (38% of poor responder in MetS patients vs. 18% in non-MetS, Odds Ratio (OR) 2.36 [Confidence Interval (CI) 1.12–6.25],  $p = 0.02$ ), while the prevalence of a MetS remained unchanged over time. The overall changes in lipid levels in our cohort under TNF blocker therapy are in line with a meta-analysis [24].

These findings suggest that a MetS may contribute to a poorer response to therapy in RA patients. This could be explained by the additive effect of the low-grade components of inflammation present in MetS to the already present inflammatory burden of rheumatoid arthritis, and partly by the pathogenetic processes driving inflammation in the two conditions (RA and MetS) that might involve different pathways and cytokines, thus requiring different therapeutic targets. The presence of a MetS could maintain some degree of chronic inflammation and, on the other hand, the persistence of inflammation associated with rheumatoid arthritis might affect the cardiovascular risk, thus mutually reinforcing each other. These data strongly suggest that it is likely to be necessary in patients with both RA and MetS that maintain a higher disease activity, compared to RA patients without MetS, to treat both conditions simultaneously and effectively in order to have a significant impact on the cardiovascular risk. This hypothesis should be tested when looking at the cardiovascular mortality rate in patients receiving DMARDs and/or biologic therapy. The introduction of new biologic therapies, that potently suppress inflammation, has further improved the outcomes but they have shown some controversial effects, especially on lipid profile. In particular, a great number of studies on anti-TNF agents have found different results, although the majority showed an increase of total cholesterol levels, as well as of HDL cholesterol, thus with an atherogenic index that tends to remain the same. Along this line, patients treated with the anti-IL6 drug, tocilizumab, are likely to experience an increase of total cholesterol, LDL, HDL and triglycerides, even though these are reversible. The long-term effect of these drugs on CV outcomes has yet to be fully elucidated [25, 26], but it is increasingly recommended to normalise cholesterol through statins, taking advantage of the intrinsic anti-inflammatory effects [27–29]. On

a large scale basis, statins do not show any clear-cut effect on RA disease activity [30], even though being on statins has shown a reduced risk of developing RA [31].

### Cardiovascular risk factors in RA are not the same as for the general population.

Hypertension, diabetes mellitus or hyperlipidemia levels among RA patients are similar to the general population. Gonzalez et al [32] found that, with the exception of smoking which also increases the susceptibility to RA, the distribution of the other traditional CV risk factors does not appear to differ between RA patients, at the time of the RA onset, and non-RA people. As the prevalence of traditional CV risk factors does not seem to account for the increased risk of CV morbidity and mortality in RA patients, the logical conclusion should be that several RA-related risk factors have to play a crucial role in the course of the disease. These factors certainly include a diminished exercise capacity, but also the possible iatrogenic effects of therapeutic interventions and chronic inflammation.

Since it is well known that disability, as measured by the Health Assessment Questionnaire (HAQ), is a predictor of both overall and cardiovascular mortality [33], the take-home message should be that HAQ remission should be included among the major outcomes for defining remission in all RA cohorts.

All the evidence supports the concept that chronic inflammation characterising RA plays a key role in accelerating atherosclerosis [34, 35]. CV disease mortality is higher in patients with more widespread disease and high levels of systemic inflammation markers [11], and Provan *et al* showed that patients with active RA, but not those in remission, had significantly increased levels of CVD risk markers (NT-proBNP, hypertension, total cholesterol, reactive hyperaemia index (RHI), measures of arterial stiffness and intima media thickness) than the control group [26]. These results indirectly support the notion that remission in RA allows diminished cardiovascular morbidity. The remaining open question is when and how we can define remission in a RA patient. It has previously been said that the increased risk of heart disease precedes the clinical onset of

rheumatoid arthritis [11], suggesting that other factors (e.g., environmental or genetic), in addition to those described above, may contribute to this early risk. In this regard, it has already been demonstrated that there is a pre-clinical phase of RA during which inflammatory activity and serological and autoimmune disturbances occur [27], thus triggering the CV risk, and that there may be a consistent delay between the onset of the first RA symptom, diagnosis and the start of an effective therapy. These findings suggest that an untreated systemic inflammation can induce damage to the CV system before it affects the joints, and most importantly that chronic exposure to systemic inflammation increases the risk of CVD [3].

The specific background of RA is clear when one considers that a lower body mass index is also associated with a higher CV mortality in RA, which is very likely related to the increased inflammatory cytokines inducing a catabolic state: the so-called rheumatoid cachexia [36].

To demonstrate that there are important and independent RA-related factors contributing to the CV risk, Solomon *et al* evaluated both traditional CV risk factors and parameters of RA severity (long disease duration, modified HAQ score, Clinical Disease Activity Index-CDAI, seropositivity, radiographic joint erosions, subcutaneous rheumatoid nodules and previous joint replacement) at baseline in predicting CV events. In this large cohort of patients the authors showed that traditional CV and RA related factors were independent predictors of CV diseases and that the risk increased with the number of both types of parameters [37].

### Metabolic syndrome and cardiovascular risk

Metabolic syndrome (MetS) represents a cluster of cardiovascular disease risk factors that have insulin resistance and increased visceral adiposity in common. This entity has received great attention as the best known predisposing setting for the development of CV morbidity [38] and represents a condition which has been defined, by the National Cholesterol Education Program's Adult Treatment Panel III Report (NCEP ATP III) from the National Institute of Health, as the situation in which three of five characterist-

**Table 1:** Behaviour over time of the metabolic variables in the 115 patients treated with a TNF $\alpha$  blocker (Adalimumab).

Baseline disease activity DAS 4.7 $\pm$ 1.1	Baseline	T6	p	T12	p
Glucose	88.1 $\pm$ 21	87.5 $\pm$ 21.6	0.370	88.3 $\pm$ 16	0.390
Glucose abn.	27(19.9%)	23(16.9%)	0.103	22(16.3%)	0.025
TGL mg/dl	122.3 $\pm$ 67.2	131.6 $\pm$ 56.3	<0.001	134.5 $\pm$ 67.7	<0.001
TGL abn	29(21.3%)	31(22.8%)	0.414	32(23.5%)	0.366
HDL mg/dl	54.0 $\pm$ 15.3	54.8 $\pm$ 13.3	0.210	55.3 $\pm$ 15.2	0.310
HDL abn	48(35.3%)	42(30.9%)	0.157	41(30.1%)	0.194
Cholesterol mg/dl	184.1 $\pm$ 37.7	191.1 $\pm$ 40.3	0.001	192.8 $\pm$ 41.3	0.002

**Table 2:** Low disease activity according to the baseline metabolic abnormalities in the cohort of patients treated with a TNF $\alpha$  blocker.

Variables	Outcome DAS $\leq$ 2.4 (54)	Outcome DAS>2.4 (61)	p
Age (years)	52 $\pm$ 12.9	55.6 $\pm$ 11.2	0.170
Sex (male)	11(20.4%)	6(9.8%)	0.120
Dis Duration (years)	8.6 $\pm$ 10.8	8.8 $\pm$ 9.3	0.640
BMI	25.5 $\pm$ 3.2	27.4 $\pm$ 4.0	0.013
BMI abnormal	23(42.6%)	39(63.9%)	0.026
MetS	10(18.5%)	23(37.7%)	0.025

ics are present, including obesity, elevated triglyceride, low level of high density lipoproteins (HDL) cholesterol, high systolic and diastolic blood pressure, and elevated fasting glucose [39]. The prevalence of MetS in the general population varies between 17 and 43 % [41, 42] and increases with age. In our cohort of RA patients receiving anti-TNF it was 27%. The MetS-associated increase in CV disease morbidity may depend on the definition used and a recent study showed that CV disease prevalence was more pronounced when the NCEP ATP III and the AHA/NHLBI criteria for MetSyn were used [42].

A recent systematic review and meta-analysis of the literature, involving more than 950'000 patients, has shown that metabolic syndrome is associated with a 2-fold increased risk of cardiovascular disease, CV mortality, myocardial infarction and stroke [43]. These data are even stronger when considering that the CV risk associated with MetS remains the same when patients with type 2 diabetes mellitus are excluded from the analyses. As a whole, MetS is a risk for higher inflammation, higher CV risk and persistence of active disease [44–49].

### Adiposity and joint inflammation

Being overweight is a major component of MetS and is associated with an adverse cardiovascular risk profile, characterised by hypertension, insulin resistance and an atherogenic lipid profile. In addition, there is a continuous relationship between BMI and risk of death from coronary artery diseases in middle-aged adults [44]. Obesity is now regarded as a systemic, low-grade inflammatory state, characterised by elevated circulating levels of C-reactive protein, TNF- $\alpha$ , IL-6 and PAI-1 [51]. Obesity is also a recognised risk factor for osteoarthritis and it is thought to be an additive risk for long standing RA [52], while there is a debate as to the possibility that obesity protects the joints in the early years of the disease [53].

Biological data have certainly clarified that adipose tissue is a dynamic endocrine organ that releases several bioactive substances, in common with inflammatory diseases such as RA, including some pro-inflammatory cytokines like TNF- $\alpha$  and IL-6, and specific cytokines, termed adipokines. These include adiponectin, leptin, resistin and visfatin, some of which promote inflammatory responses and metabolic dysfunction, and others which contribute to the resolution of inflammation and have beneficial effects on obesity-linked metabolic disorders [54]. One adipokine which gained utmost importance just recently has been chemerin, which plays a critical role in adipogenesis. Through its receptor ChemR23 it can amplify inflammation and lead to cartilage damage. In obese people it has been shown that adipose tissue presents more inflammatory infiltrate than normal-weight people and is characterised by different regulatory mechanisms and cytokine pattern production [55]. This implies that pro-inflammatory cytokines and adipokines can affect metabolic dysfunction and CV risk on the one hand, and rheumatoid arthritis on the other one. In this regard, an Italian retrospective study first described an approximately 3-fold lower chance to achieve remission in RA, obese patients receiving anti-TNF therapy than in normal-weight RA patients [56]. These data were

confirmed by a recent prospective study showing that a high body mass index (BMI) was associated with a poor response to infliximab [57], a drug whose dose should still be proportional to body weight, supporting the fact that adipose tissue may be involved in the pathophysiology of RA. If we consider that obese patients are at increased risk of developing RA, with an adjusted odds ratio of 3.74 [58], then we may conclude that an obese RA patient does have more inflammation than the same RA in a non-obese patient.

### Adiposity, adipo-cytokines, joints and vessels

#### TNF-alpha

In white adipose tissue (WAT), TNF- $\alpha$  is produced by adipocytes and fat infiltrating macrophages and is over-expressed in plasma and adipose tissue of obese human and animal models. TNF- $\alpha$  promotes insulin resistance by decreasing tyrosine kinase activity of the insulin receptor and insulin signalling *via* MAPK pathways *in vitro* and *in vivo*, thereby reducing insulin activity [59, 60]. Studies in mice suggest that the attenuation of TNF activity improves glucose homeostasis [61]. In humans in a small group of obese T2DM patients, a study showed that blocking TNF- $\alpha$  decreased plasma levels of inflammatory markers but did not ameliorate insulin resistance [62], while a recent report found that a prolonged period of TNF neutralisation in patients with a metabolic syndrome improved fasting glucose levels and increased high molecular weight adiponectin levels [63]. Moreover, anti-TNF therapy in “high-grade” inflammatory diseases, such as rheumatoid arthritis, has been reported to promote insulin sensitivity in several small studies [64, 65], suggesting that TNF has an important role in inducing insulin resistance mediated by inflammation.

TNF- $\alpha$  also plays a role in endothelial dysfunction [4, 66], an early marker of atherosclerosis, and as such is already




Figure 1

Fat inflammation increases joint inflammation and cardiovascular risk.

present in MetS [67]. There is evidence that TNF- $\alpha$  blockade in RA patients leads to a decrease of biomarkers of endothelial dysfunction and an improvement of endothelial dysfunction [68, 69], especially in the early phases of the disease [4].

### IL-6

WAT produces 10 to 30% of circulating IL6, and plasma levels of IL-6 strongly correlate with BMI and insulin resistance [70]. Elevated IL-6 plasma levels predict future risk of cardiovascular events [71] and of T2DM [72], independently of BMI.

In a preliminary study involving a small group of non diabetic RA patients, it seemed that inhibition of IL-6 signalling improved insulin sensitivity [73]. The IL-6 effects on accelerated atherosclerosis are controversial. In an experimental model, administration of exogenous IL-6 in the murine apolipoprotein E-deficient (*ApoE*<sup>-/-</sup>) model of atherosclerosis dramatically enhanced atherosclerotic lesion formation, suggesting a pivotal role for IL-6 in plaque progression [74]. However, a lifetime deficiency of IL-6 in the *ApoE*<sup>-/-</sup> model enhanced atherosclerotic lesion formation and increased cholesterol levels [75, 76].

In light of these data, it will be interesting to see if antagonising IL-6 in RA positively affects the cardiovascular outcomes, despite the known negative effects on the lipid profile. These data will certainly derive from inception cohorts in registries.

### Leptin

WAT also produces the adipokine leptin, the product of the *ob* gene [77]. Leptin is considered a major regulator of body weight by suppressing appetite, thus decreasing food intake, and stimulating energy expenditure via hypothalamic receptors. Plasma levels of leptin are directly correlated with the amount of body fat and obese people have high circulating levels of leptin, indicating the occurrence of resistance to its action [78]. However, leptin is also involved in the inflammatory and immune mechanisms. Leptin synthesis is enhanced during acute infection and inflammation [79], as well as under the action of inflammatory mediators such as TNF $\alpha$ , IL-6 and IL-1. Furthermore, leptin stimulates the production of pro-inflammatory cytokines from cultured monocytes and increases the production of Th1 type cytokines, down-regulating Th2-type cytokines, thus polarising T cells towards a Th1 phenotype [80–81]. High circulating leptin levels have recently been demonstrated in patients with RA [82], both in plasma and in the synovial fluid. The clinical relevance of these findings still needs to be fully elucidated [83, 84].

### Resistin

Resistin is a 108 amino acid long protein (12.5 kD) initially related to insulin resistance [85]. In animal models, resistin promotes insulin resistance, while the evidence for this effect in humans is less clear [86, 87]. Resistin was originally identified in adipose tissue. Later, it was observed that resistin production is restricted to adipocytes in mice, while in humans it is mainly derived from circulating monocytes and macrophages [88]. In this regard, several studies demonstrated that pro-inflammatory cytokines increase the

expression of resistin [89], which in turn is able to stimulate the production of TNF $\alpha$  and IL12 [90], and to up-regulate the genes of TNF- $\alpha$ , IL-6, IL-1 $\beta$  and, interestingly, resistin itself [91]. Being produced by macrophages, resistin appears to be one of the several enhancers of inflammation produced by these cells. Along this line, it has been suggested that resistin may have a role in the pathogenesis of arthritis, triggering inflammatory synovitis when injected into mice joints [92]. In rheumatoid arthritis, despite conflicting results concerning serum levels, resistin synovial fluid levels were found to be higher in RA when compared to osteoarthritis, and resistin expression in synovial lining layers positively correlated with acute phase reactants and disease activity [93]. Of interest, recent experimental data suggest that resistin, in the presence of dendritic cells (DC), might induce the expansion of functional regulatory T cells (TRegs) [94].

### Adiponectin

Adiponectin, encoded by the ADIPOQ gene, is a protein which is almost exclusively synthesised and secreted by adipocytes. It shows anti-inflammatory, insulin-sensitising and anti-atherogenic properties [95]. Adiponectin plasma levels are decreased in obese people compared with normal weight individuals, and also reduced in type 2 diabetes and MetS. These findings suggest that the functional adipocytes of lean subjects express high adiponectin levels, while in the dysfunctional adipocytes of obese individuals its expression is down-regulated. Moreover, weight reduction significantly increases circulating levels.

Adiponectin has shown several “anti-inflammatory” effects: it inhibits the transformation of macrophages into foam cells; it reduces TNF production by macrophages; it promotes the alternative activation of monocytes from a pro-inflammatory M1-type macrophage into an anti-inflammatory M2-phenotype; and it stimulates the production of the anti-inflammatory cytokine IL10 and of Interleukin 1 Receptor antagonist (IL1Ra) by macrophages [96, 97]. Studies in RA found elevated serum and synovial fluid adiponectin levels both in early and established disease, which did not correlate with disease activity nor were affected by TNF blockade, but increased with methotrexate treatment [98]. These data suggest that Adiponectin induces a sort of “natural” tolerance in macrophages to inflammatory stimuli.

### Visfatin

Visfatin is a novel 55 kD adipokine that was originally discovered in the liver, skeletal muscle and bone marrow as a growth factor for B lymphocyte precursors (hence pre-B-cell colony-enhancing factor 1, PBEF) with insulin-mimetic actions, preferentially produced by visceral adipose tissue but also found in the liver, bone marrow, skeletal muscle and lymphocytes [99, 100]. Circulating levels of visfatin correlate with obesity and type-2 diabetes, and are reduced after weight loss [101].

Visfatin is also synthesised in response to inflammatory stimuli, and, moreover, it can enhance production of TNF- $\alpha$ , IL-6, IL-1 $\beta$ , as well as of IL-10 and IL-1Ra [102], and could promote atherosclerotic processes (inducing cell adhesion molecules) and plaque destabilisation [103]. Visfatin

has been shown to induce chemotaxis and the production of IL-1 $\beta$ , TNF, IL-6 and co-stimulatory molecules by CD14<sup>+</sup> monocytes, and to increase their ability to induce all proliferative responses in lymphocytes, effects which are mediated intracellularly by p38 and MEK1 [104]. Inhibiting visfatin reduced circulating TNF $\alpha$  levels during endotoxemia in mice [105] in other inflammatory conditions, and also in rheumatoid arthritis some authors reported increased synovial fluid and plasma levels of this adipokine as well as its expression by rheumatoid synoviocytes at sites of attachment and invasion into cartilage or bone [106]. Additionally in agreement with these findings is the association between radiographic damage and high visfatin serum concentrations [107].

#### BAFF (B cell activating factor)

BAFF is a 17 kD protein mainly secreted by myeloid cells (but also by bone marrow stromal cells, granulocytes, synovial fibroblasts etc.) that exerts profound effects on B cells leading to their maturation and increased survival. BAFF has three receptors: B cell maturation antigen (BCMA), trans-membrane activator and CAML interactor (TACI), and BAFF receptor (BAFF-R) [108; 109].

It has recently been shown that BAFF and its receptors are synthesised by maturing adipocytes, that TNF increased BAFF synthesis by mature adipocytes, and that BAFF itself increased the synthesis of two of its receptors, namely BAFF-R and BCMA [110]. Whether the expression of the ligand and of its receptors might influence the course or the occurrence of autoimmune disease will only be solved by prospective studies.

#### Chemerin and ChemR23

Chemerin is a novel 18 kD adipokine that regulates adipocyte development and metabolic function as well as glucose metabolism in liver and skeletal muscle tissues. The chemerin gene, also known as tazarotene-induced gene 2 (TIG2) or retinoic acid receptor responder 2 (RARRES2), was originally identified as a novel retinoid-responsive gene in psoriatic skin lesions. Chemerin is a secreted protein that exerts its functions by binding the G protein-coupled receptor ChemR23. It was first discovered as a chemotactic peptide directing macrophages and dendritic cells expressing ChemR23 towards sites of inflammation, being involved in both adaptive and innate immunity [111, 112].

ChemR23 and chemerin proteins have been detected in chondrocytes *in vitro* by immunocytochemistry, and ChemR23 and prochemerin transcripts were detected in chondrocyte cultures by reverse transcriptase PCR. It has been shown that Chemerin binding to ChemR23 led to increased phosphorylation of p44/42 mitogen-activated protein kinases (MAPKs) and Akt and blocking of MEK-1/2 signalling prevented phosphorylation of p44/42 MAPKs, but not of Akt. This suggests that intracellular downstream events upon chemerin stimulation occur through the Akt/MEK/MAPK pathway [113]. In chondrocyte cultures, Chemerin regulates the production of pro-inflammatory cytokines, IL-1 $\beta$ , TNF- $\alpha$ , IL-6, and IL-8 as well as MMP-13, thus suggesting that it can lead to cartilage damage and degradation [114]. Indirect evidence for this hypothesis

comes from the observation that Chemerin levels are increased in synovial fluid of RA patients and that synovial tissue from RA shows expression of both Chemerin and ChemR23 [115]. Therefore, obesity and joint damage seems to be linked by several possible pathways (Fig. 1). The interesting aspect of this possible pathway is that Chemerin-derived peptides, after proteolysis of the precursor protein, can also exert anti-inflammatory effects, thus suggesting that the initial synthesis and the chronic production of Chemerin can have different effects [116]. Whether this might explain why obesity in the early phases of RA seems to exert a protective effect whereas it does have a deleterious effect over the long lasting periods, will be solved by dedicated prospective studies.

#### Conclusions

In view of the role of adipokines in inflammatory arthritis and the potential modulatory role of TNF on adipokines, some studies have tested the effect of TNF blockade on the plasma levels of some of these adipokines in patients with RA. In particular, with regard to adiponectin, five studies have found that short or long-term TNF blockade had no influence on circulating levels of this cytokine [117], while four showed an increase of adiponectin after anti-TNF administration [118–119]. Additionally, four studies on leptin [120] and one on visfatin [121] showed no changes in plasma concentrations of these adipokines and one study described a decrease of resistin following anti-TNF therapy [122–124]. Only one study showed that a TNF $\alpha$  blocker decreased chemerin levels [115].

The conclusion at this moment is that the biological role of adipokines is not entirely understood. The evidence that obesity represents a negative biomarker for TNF $\alpha$  blockers-treated patients to reach low disease activity or remission, suggests that it represents an amplifying loop on the course of RA. Only an adequate control of obesity and MetS seems to guarantee full control of the systemic inflammation and, as such, of the related CV risk.

**Funding / potential competing interests:** Supported by ASRALES foundation.

**Correspondence:** Professor Gianfranco Ferraccioli, MD, Division of Rheumatology, Catholic University of the Sacred Heart, Via Moscatti 31, IT-00168, Rom, Italy, [gf.ferraccioli@rm.unicatt.it](mailto:gf.ferraccioli@rm.unicatt.it)

#### References

- Avina-Zubieta JA, Choi HK, Sadatsafavi M, Etminan M, Esdaile JM, Lacaille D. Risk of cardiovascular mortality in patients with rheumatoid arthritis: a meta-analysis of observational studies. *Arthritis Rheum* 2008;59:1690–7.
- Gabriel SE. Heart disease and rheumatoid arthritis: understanding the risk. *Ann Rheum Dis* 2010;69(Suppl):i61–i64.
- Ku IA, Imboden JB, Hsue PY, Ganz P. Rheumatoid arthritis: model of systemic inflammation driving atherosclerosis. *Circ J* 2009;73:977–85.
- Cardillo C, Schinzari F, Mores N, Mettimano M, Melina D, Zoli A, Ferraccioli G. Intravascular tumor necrosis factor alpha blockade reverses

- endothelial dysfunction in rheumatoid arthritis. *Clin Pharmacol Ther.* 2006;80:275–81.
- 5 Watson DJ, Rhodes T, Guess HA. All-cause mortality and vascular events among patients with rheumatoid arthritis, osteoarthritis, or no arthritis in the UK General Practice Research Database. *J Rheumatol* 2003;30: 1196–1202.
- 6 Wolfe F, Mitchell DM, Sibley JT, Fries JF, Bloch DA, Williams CA, et al. The mortality of rheumatoid arthritis. *Arthritis Rheum* 1994;37:481–494.
- 7 Naranjo A, Sokka T, Descalzo MA, Calvo-Alén J, Hørslev-Petersen K, Luukkainen RK, et al; QUEST-RA Group. Cardiovascular disease in patients with rheumatoid arthritis: results from the QUEST-RA study. *Arthritis Res Ther* 2008;10:R30.
- 8 van Halm VP, Peters MJ, Voskuyl AE, Boers M, Lems WF, Visser M, et al. Rheumatoid arthritis versus diabetes as a risk factor for cardiovascular disease: a cross-sectional study, the CARRÉ Investigation. *Ann Rheum Dis* 2009;68:1395–400.
- 9 Peters MJ, van Halm VP, Voskuyl AE, Smulders YM, Boers M, Lems WF, et al. Does rheumatoid arthritis equal diabetes mellitus as an independent risk factor for cardiovascular disease? A prospective study. *Arthritis Rheum* 2009;61:1571–9.
- 10 Ferraccioli G, Gremese E. Autoantibodies and thrombophilia in RA: TNF $\alpha$  and TNF $\alpha$  blockers. *Ann Rheum Dis* 2004;63:613–15.
- 11 Maradit-Kremers H, Crowson CS, Nicola PJ, Ballman KV, Roger VL, Jacobsen SJ, et al. Increased unrecognized coronary heart disease and sudden deaths in rheumatoid arthritis: a population-based cohort study. *Arthritis Rheum.* 2005;52:402–11.
- 12 Gabriel SE. Cardiovascular morbidity and mortality in rheumatoid arthritis. *Am J Med.* 2008;121:S9–14.
- 13 Aubry MC, Maradit-Kremers H, Reinalda MS, Crowson CS, Edwards WD, Gabriel SE. Differences in atherosclerotic coronary heart disease between subjects with and without rheumatoid arthritis. *J Rheumatol.* 2007;34:937–42.
- 14 Gonzalez A, Maradit-Kremers H, Crowson CS, Nicola PJ, Davis JM, Therneau TM, et al. The widening mortality gap between rheumatoid arthritis patients and the general population. *Arthritis Rheum* 2007; 56: 3583 – 3587.
- 15 Kremers HM, Crowson CS, Therneau TM, Roger VL, Gabriel SE. High ten-year risk of cardiovascular disease in newly diagnosed rheumatoid arthritis patients: a population-based cohort study. *Arthritis Rheum* 2008;58:2268–74.
- 16 Meune C, Touzé E, Trinquart L, Allanore Y. Trends in cardiovascular mortality in patients with rheumatoid arthritis over 50 years: a systematic review and meta-analysis of cohort studies. *Rheumatology* 2009;48:1309–13.
- 17 Peltomaa R, Paimela L, Kautiainen H, Leirisalo-Repo M. Mortality in patients with rheumatoid arthritis treated actively from the time of diagnosis. *Ann Rheum Dis* 2002;61:889–94.
- 18 Carmona L, Descalzo MA, Perez-Pampin E, Ruiz-Montesinos D, Erra A, Cobo T, et al, and the BIOBADASER and EMECAR Groups. All-cause and cause-specific mortality in rheumatoid arthritis are not greater than expected when treated with tumor necrosis factor antagonists. *Ann Rheum Dis* 2007;66:880–5.
- 19 Greenberg JD, Kremer JM, Curtis JR, Hochberg MC, Reed G, Tsao P, et al; on behalf of the CORRONA Investigators. Tumor necrosis factor antagonist use and associated risk reduction of cardiovascular events among patients with rheumatoid arthritis. *Ann Rheum Dis.* 2010 Nov 24. Epub ahead of print
- 20 Al-Aly Z, Pan H, Zeringue A, Xian H, McDonald JR, El-Achkar TM, Eisen S. Tumor necrosis factor- $\alpha$  blockade, cardiovascular outcomes, and survival in rheumatoid arthritis. *Transl Res* 2011;157:10–8.
- 21 Pereira RM, de Carvalho JF, Bonfá E. Metabolic syndrome in rheumatological diseases. *Autoimmun Rev* 2009;8:415–9.
- 22 Karvounaris SA, Sidiropoulos PI, Papadakis JA, Spanakis EK, Bertias GK, Kritikos HD et al. Metabolic syndrome is common among middle-to-older-age Mediterranean patients with rheumatoid arthritis and correlates with disease activity: a retrospective, cross-sectional, controlled study. *Ann Rheum Dis* 2007;66:28–33.
- 23 Chung CP, Oeser A, Solus JF, Avalos I, Gebretsadik T, Shintani A, et al. Prevalence of the metabolic syndrome is increased in rheumatoid arthritis and is associated with coronary atherosclerosis. *Atherosclerosis* 2008;196:756–63.
- 24 Steiner G, Urowitz M.B. Lipid profiles in patients with rheumatoid arthritis: mechanisms and the impact of treatment. *Semin Arthritis Rheum.* 2009;38:372–81
- 25 Ferraccioli G, Gremese E. Thrombogenicity of TNF alpha in rheumatoid arthritis defined through biological probes: TNF alpha blockers. *Autoimmun Rev* 2004 Jun;3:261–6.
- 26 Provan SA, Semb AG, Hisdal J, Strandén E, Agewall S, Dagfinrud H, et al. Remission is the goal for cardiovascular risk management in patients with rheumatoid arthritis: a cross-sectional comparative study. *Ann Rheum Dis* 2011 Feb 2 Epub ahead of print.
- 27 Kokkonen H, Soderstrom I, Rocklov J, Hallmans G, Lejon K, Rantapaa-Dalqvist S. Up-regulation of cytokines and chemokines predates the onset of rheumatoid arthritis. *Arthritis Rheum* 2010;62:383–91.
- 28 Peters MJ, Symmons DP, McCarey D, Dijkman BA, Nicola P, Kvien TK, et al. EULAR evidence-based recommendations for cardiovascular risk management in patients with rheumatoid arthritis and other forms of inflammatory arthritis. *Ann Rheum Dis.* 2010;69:325–31
- 29 Jain MK, Ridker PM: Anti-inflammatory effects of statins: clinical evidence and basic mechanisms. *Nature Reviews Drug Discovery* 4, 977–987.
- 30 Lodi S, Evans SJ, Egger P, Carpenter J. Is there an anti-inflammatory effect of statins in rheumatoid arthritis? Analysis of a large routinely collected claims database. *Br J Clin Pharmacol.* 2010;69:85–94.
- 31 Chodick G, Amital H, Shalem Y, Kokia E, Heymann AD, Porath A, Shalev V. Persistence with statins and onset of rheumatoid arthritis: a population-based cohort study. *PLoS Med.* 2010;7:e1000336.
- 32 Gonzalez A, Maradit-Kremers H, Crowson CS, Ballman KV, Roger VL, Jacobsen SJ, et al. Do cardiovascular risk factors confer the same risk for cardiovascular outcomes in rheumatoid arthritis patients as in non-rheumatoid arthritis patients? *Ann Rheum Dis* 2008;67:64–9.
- 33 Turesson C, Matteson EL. Cardiovascular risk factors, fitness and physical activity in rheumatic diseases. *Curr Opin Rheumatol* 2007;19:190–6.
- 34 Gremese E, Ferraccioli G. The metabolic syndrome: the crossroads between rheumatoid arthritis and cardiovascular risk. *Autoimmun Rev* 2011 ahead of print.
- 35 Abou-Raya A, Abou-Raya S. Inflammation: a pivotal link between autoimmune diseases and atherosclerosis. *Autoimmun Rev* 2006;5:331–7.
- 36 Escalante A, Haas RW, del Rincon I. Paradoxical effect of body mass index on survival in rheumatoid arthritis: Role of comorbidity and systemic inflammation. *Arch Intern Med* 2005;165:1624–1629.
- 37 Solomon DH, Kremer J, Curtis JR, Hochberg MC, Reed G, Tsao P, et al. Explaining the cardiovascular risk associated with rheumatoid arthritis: traditional risk factors versus markers of rheumatoid arthritis severity. *Ann Rheum Dis* 2010;69:1920–5.
- 38 Hu G, Qiao Q, Tuomilehto J, Balkau B, Borch-Johnsen K, Pyorala K. Prevalence of the metabolic syndrome and its relation to all cause and cardiovascular mortality in nondiabetic European men and women. *Arch Int Med* 2004;164:1066–76.
- 39 National Institutes of Health, 2001. Third report of the National Cholesterol Education Program Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). National Institutes of Health, Bethesda, Md. NIH Publication 01–3670.
- 40 Ford ES, Giles WH, Dietz WH. Prevalence of the metabolic syndrome among US adults. Findings from the third National Health and Nutrition Examination Survey. *JAMA* 2002;287:356–359.
- 41 Bonora E, Kiechl S, Willeit J, Oberlorenz F, Egger G, Bonadonna RC et al. Metabolic Syndrome: epidemiology and more extensive phenotypic description. Cross-sectional data from the Bruneck Study. *Int J Obes Relat Metab Disord* 2003;27:1283–89.
- 42 Athyros VG, Ganotakis ES, Elisaf MS, Liberopoulos EN, Goudevenos IA, Karagiannis A; for the GREECE-METS Collaborative Group. Pre-

- valence of vascular disease in metabolic syndrome using three proposed definitions. *Int J Cardiol* 2007;117:204–10.
- 43 Mottillo S, Filion KB, Genest J, Joseph L, Pilote L, Poirier P, et al. The metabolic syndrome and cardiovascular risk a systematic review and meta-analysis. *J Am Coll Cardiol* 2010;56:1113–32.
- 44 Dessein PH, Tobias M, Veller MG: Metabolic syndrome and subclinical atherosclerosis in rheumatoid arthritis. *J Rheumatol* 2006;33:2425–32
- 45 Zonana-Nacach A, Santana-Sahagun E, Jimenez-Balderas FJ, Camargo-Coronel A: Prevalence and factors associated with metabolic syndrome in patients with RA and SLE: *J Clin Rheumatol* 2008;14:74–7
- 46 Toms TE, Panoulas VF, John H, Douglas KM, Kitas GD: Methotrexate therapy associates with reduced prevalence of the metabolic syndrome in RA patients over age of 60—more than just an anti-inflammatory effect? a cross sectional study: *Arthritis Res Ther* 2009;11:R110
- 47 Elkan AC, Hakansson N, Frostegard J, Cederholm T, Brismar K, Hafstrom I: Cachexia in rheumatoid arthritis is associated with inflammatory activity, physical disability and low bioavailable insulin-like growth factor. *Arthritis Res Ther* 2009;11:R37
- 48 Ratterman HG, Van Eijk IC, Voskuil AE, Peters MJ, Dijkmans BA, Van Halm VP, Simsek S, Lems WF, Nurmohamed MT: The metabolic syndrome is amplified in hypothyroid rheumatoid arthritis patients: a cross sectional study. *Ann Rheum Dis* 2010;69:39–42
- 49 Crowson CS, Myasoedova E, Davis JM, Matteson EL, Roger VL, Therneau TM, Fitz-Gibbon P, Rodeheffer RJ, Gabriel SE: Increased prevalence of metabolic syndrome associated with rheumatoid arthritis in patients without clinical cardiovascular disease: *J Rheumatol* 2011, 38: 29–35
- 50 Dao HH, Do QT, Sakamoto J: Increased frequency of metabolic syndrome among Vietnamese women with early rheumatoid arthritis: a cross sectional study: *Arthritis Res Ther* 2010;12:R218
- 51 Prospective Studies Collaborations. Body-mass index and cause-specific mortality in 900 000 adults: collaborative analyses of 57 prospective studies. *Lancet* 2009;373:1083–1096.
- 52 Stavropoulos-Kalinoglou A, Metsios GS, Koutedakis Y, Kitas GD. Obesity in rheumatoid arthritis. *Rheumatology (Oxford)*. 2011;50:450–62
- 53 Westhoff G, Rau R, Zink A. Radiographic joint damage in early rheumatoid arthritis is highly dependent on body mass index. *Arthritis Rheum*. 2007;56:3575–82
- 54 Ouchi N, Parker JL, Lugus JJ, Walsh K. Adipokines in inflammation and metabolic disease. *Nat Rev Immunol* 2011;11:85–97.
- 55 Sopasakis VR, Nagaev I, Smith U. Cytokine release from adipose tissue of nonobese individuals. *Int J Obes* 2005; 29: 1144–1147
- 56 Ferraccioli G, Trotta F, Punzi L, Ferri C, Sarzi Puttini PC, Bambara LM, et al. Body weight and response to biologicals in rheumatoid arthritis and Spondylarthritis. Obesity reduces the rate of remission-response. The GISEA registry. Abstract The Annual European Congress of Rheumatology. Italy, Rome June 2010.
- 57 Klaasen R, Wijbrandts CA, Gerlag DM, Tak PP. Body mass index and clinical response to infliximab in rheumatoid arthritis. *Arthritis Rheum* 2011;63:359–64.
- 58 Symmons DP, Bankhead CR, Harrison BJ, Brennan P, Barrett EM, Scott DG, et al. Blood transfusion, smoking, and obesity as risk factors for the development of rheumatoid arthritis: results from a primary care-based incident case-control study in Norfolk, England. *Arthritis Rheum* 1997;40:1955–61.
- 59 Hotamisligil GS, Budavari A, Murray D, Spiegelman BM. Reduced tyrosine kinase activity of the insulin receptor in obesity-diabetes. Central role of tumor necrosis factor-alpha. *J Clin Invest* 1994;94:1543–9.
- 60 Hotamisligil GS, Peraldi P, Budavari A, Ellis R, White MF, Spiegelman BM. IRS-1-mediated inhibition of insulin receptor tyrosine kinase activity in TNF-alpha- and obesity-induced insulin resistance. *Science* 1996;271:665–8.
- 61 Uysal KT, Wiesbrock SM, Marino MW, Hotamisligil GS. Protection from obesity-induced insulin resistance in mice lacking TNF-alpha function. *Nature* 1997;389:610–614.
- 62 Dominguez H, Storgaard H, Rask-Madsen C, Steffen Hermann T, Ihlemann N, Baunbjerg Nielsen D, et al. Metabolic and vascular effects of tumor necrosis factor-alpha blockade with etanercept in obese patients with type 2 diabetes. *J Vasc Res* 2005;42:517–25.
- 63 Stanley TL, Zanni MV, Johnsen S, Rasheed S, Makimura, Lee H, et al. TNF-alpha antagonism with Etanercept decreases glucose and increases the proportion of high molecular weight adiponectin in obese subjects with features of the metabolic syndrome. *J Clin Endocrinol Metab* 2011;96:E146–50.
- 64 Gonzalez-Gay MA, Gonzalez-Juanatey C, Vazquez-Rodriguez TR, Miranda-Filloo JA, Llorca J. Insulin resistance in rheumatoid arthritis: the impact of the anti-TNF-alpha therapy. *Ann N Y Acad Sci* 2010;1193:153–9.
- 65 Ursini F, Naty S, Grembiale RD. Infliximab and insulin resistance. *Autoimmun Rev* 2010;9:536–9.
- 66 Chia S, Qadan M, Newton R, Ludlam CA, Fox KA, Newby DE. Intra-arterial tumor necrosis factor-alpha impairs endothelium-dependent vasodilatation and stimulates local tissue plasminogen activator release in humans. *Arterioscler Thromb Vasc Biol* 2003;23:695–701.
- 67 Tziomalos K, Athyros VG, Karagiannis A, Mikhailidis DP. Endothelial dysfunction in metabolic syndrome: prevalence, pathogenesis and management. *Nutr Metab Cardiovasc Dis*. 2010;20:140–6.
- 68 Hürlimann D, Forster A, Noll G, Enseleit F, Chenevard R, Distler O, et al. Anti-tumor necrosis factor-alpha treatment improves endothelial function in patients with rheumatoid arthritis. *Circulation* 2002;106:2184–7.
- 69 Gonzalez-Juanatey C, Testa A, Garcia-Castelo A, Garcia-Porrúa C, Llorca J, Gonzalez-Gay MA. Active but transient improvement of endothelial function in rheumatoid arthritis patients undergoing long-term treatment with anti-tumor necrosis factor alpha antibody. *Arthritis Rheum*. 2004;51:447–50.
- 70 Fried SK, Bunkin DA, Greenberg AS. Omental and subcutaneous adipose tissues of obese subjects release interleukin-6: depot difference and regulation by glucocorticoid. *J Clin Endocrinol Metab* 1998;83:847–850.
- 71 Nishida H, Horio T, Suzuki Y, Iwashima Y, Tokudome T, Yoshihara F, et al. Interleukin-6 as an independent predictor of future cardiovascular events in high-risk Japanese patients: Comparison with C-reactive protein. *Cytokine* 2011;53:342–6.
- 72 Senn JJ, Klover PJ, Nowak IA, Mooney RA. Interleukin-6 induces cellular insulin resistance in hepatocytes. *Diabetes* 2002;51:3391–3399.
- 73 Matthews VB, Allen TL, Risis S, Chan MH, Henstridge DC, Watson N, et al. Interleukin-6-deficient mice develop hepatic inflammation and systemic insulin resistance. *Diabetologia* 2010;53:2431–41.
- 74 Schultz O, Oberhauser F, Saech J, Rubbert-Roth A, Hahn M, Krone W, et al. Effects of inhibition of interleukin-6 signalling on insulin sensitivity and lipoprotein (a) levels in human subjects with rheumatoid diseases. *PLoS One* 2010;5:e14328.
- 75 Huber SA, Sakkinen P, Conze D, Hardin N, Tracy R. Interleukin-6 exacerbates early atherosclerosis in mice. *Arterioscler Thromb Vasc Biol*. 1999;19:2364–7.
- 76 Schieffer B, Selle T, Hilfiker A, Hilfiker-Kleiner D, Grote K, Tietge UJ, et al. Impact of interleukin-6 on plaque development and morphology in experimental atherosclerosis. *Circulation* 2004;110:3493–500.
- 77 Zhang Y, Proenca R, Maffei M, Barone M, Leopold L, Friedman JM. Positional cloning of the mouse obese gene and its human homologue. *Nature* 1994;372:425–432.
- 78 Jéquier E. Leptin signaling, adiposity, and energy balance. *Ann N Y Acad Sci* 2002; 967:379–388.
- 79 Otero M, Lago R, Lago F, Casanueva FF, Dieguez C, Gómez-Reino JJ, et al. Leptin, from fat to inflammation: old questions and new insights. *FEBS Lett* 2005;579:295–301.
- 80 Montecucco F, Bianchi G, Gnerre P, Bertolotto M, Dallegrì F, Ottonello L. Induction of neutrophil chemotaxis by leptin: crucial role for p38 and Src kinases. *Ann N Y Acad Sci* 2006;1069:463–471.
- 81 Otero M, Lago R, Gomez R, Dieguez C, Lago F, Gómez-Reino J et al. Towards a pro-inflammatory and immunomodulatory emerging role of leptin. *Rheumatology* 2006;45:944–950.
- 82 Lord GM, Matarese G, Howard JK, Baker RJ, Bloom SR, Lechler RI. Leptin modulates the T-cell immune response and reverses starvation-induced immunosuppression. *Nature* 1998;394: 897–901.



- 83 Otero M, Lago R, Gomez R, Lago F, Dieguez C, Gómez-Reino JJ, et al. Changes in fat-derived hormones plasma concentrations: adiponectin, leptin, resistin, and visfatin in rheumatoid arthritis subjects. *Ann Rheum Dis* 2006;65:1198–1201.
- 84 Seven A, Güzel S, Aslan M, Hamuryudan V. Serum and synovial fluid leptin levels and markers of inflammation in rheumatoid arthritis. *Rheumatol Int* 2009;29:743–747.
- 85 Rho YH, Solus J, Sokka T, Oeser A, Chung CP, Gebretsadik T, et al. Adipocytokines are associated with radiographic joint damage in rheumatoid arthritis. *Arthritis Rheum* 2009;60:1906–1914.
- 86 Steppan CM, Bailey ST, Bhat S, Brown EJ, Banerjee RR, Wright CM, et al. The hormone resistin links obesity to diabetes. *Nature* 2001;409:307–312.
- 87 Heilbronn LK, Rood J, Janderoova L, Albu JB, Kelley DE, Ravussin E, et al. Relationship between serum resistin concentrations and insulin resistance in nonobese, obese, and obese diabetic subjects. *J Clin Endocrinol Metab* 2004;89:1844–1848.
- 88 Lee JH, Chan JL, Yiannakouris N, Kontogianni M, Estrada E, Seip R, et al. Circulating resistin levels are not associated with obesity or insulin resistance in humans and are not regulated by fasting or leptin administration: cross-sectional and interventional studies in normal, insulin-resistant, and diabetic subjects. *J Clin Endocrinol Metab* 2003;88:4848–4856.
- 89 Savage DB, Sewter CP, Klenk ES, Segal DG, Vidal-Puig A, Considine RV, et al. Resistin/Fizz3 expression in relation to obesity and peroxisome proliferator activated receptor- $\gamma$  action in humans. *Diabetes* 2001;50:2199–2202.
- 90 Kaser S, Kaser A, Sandhofer A, Ebenbichler CF, Tilg H, Patsch JR. Resistin messenger-RNA expression is increased by proinflammatory cytokines in vitro. *Biochem Biophys Res Commun* 2003;309:286–290.
- 91 Silswal N, Singh AK, Aruna B, Mukhopadhyay S, Ghosh S, Ehtesham NZ. Human resistin stimulates the pro-inflammatory cytokines TNF- $\alpha$  and IL-12 in macrophages by NF- $\kappa$ B-dependent pathway. *Biochem Biophys Res Commun* 2005;34:1092–1101.
- 92 Bokarewa M, Nagaev I, Dahlberg L, Smith U, Tarkowski A. Resistin, an adipokine with potent proinflammatory properties. *J Immunol* 2005;74:5789–5795.
- 93 Senolt L, Housa D, Vernerová Z, Jirasek T, Svobodova R, Veigl D, et al. Resistin in rheumatoid arthritis synovial tissue, synovial fluid and serum. *Ann Rheum Dis* 2007;66:458–463.
- 94 Son YM, Ahn SM, Kim GR, Moon YS, Kim SH, Park YM, et al. Resistin enhances the expansion of regulatory T cells through modulation of dendritic cells. *BMC Immunol*. 2010, 30;11:33.
- 95 Schäffler A, Ehling A, Neumann E, Herfarth H, Tarner I, Schölmerich J, et al. Adipocytokines in synovial fluid. *JAMA* 2003;290:1709–1710.
- 96 Peters MJ, Watt PH, Cherry L, Welsh P, Henninger E, Dijkmans BA, et al. Lack of effect of TNF $\alpha$  blockade therapy on circulating adiponectin levels in patients with autoimmune disease: results from two independent prospective studies. *Ann Rheum Dis* 2010;69:1687–1690.
- 97 Tsatsanis C, Zacharioudaki V, Androulidaki A, Dermizaki E, Charalampopoulos I et al. Adiponectin induces TNF and IL6 in macrophages and promotes tolerance to itself and other pro-inflammatory stimuli. *Biochem.Biophys.Res.Comm.* 2005, 335: 1254–63
- 98 Popa C, Netea MG, de Graaf J, van den Hoogen FH, Radstake TR, Toenhake-Dijkstra H, et al. Circulating leptin and adiponectin concentrations during tumor necrosis factor blockade in patients with active rheumatoid arthritis. *J Rheumatol* 2009;36:724–30.
- 99 Gonzalez-Gay MA, Vazquez-Rodriguez TR, Garcia-Unzueta MT, Berja A, Miranda-Fillooy JA, de Matias JM, et al. Visfatin is not associated with inflammation or metabolic syndrome in patients with severe rheumatoid arthritis undergoing anti-TNF- $\alpha$  therapy. *Clin Exp Rheumatol* 2010;28:56–62.
- 100 Sethi JK, Vidal-Puig A. Visfatin: the missing link between intra-abdominal obesity and diabetes? *Trends Mol Med* 2005;11:344–347.
- 101 Haider DG, Schindler K, Schaller G, Prager G, Wolzt M, Ludvik B. Increased plasma visfatin concentrations in morbidly obese subjects are reduced after gastric banding. *J Clin Endocrinol Metab* 2006;91:1578–81.
- 102 Moschen AR, Kaser A, Enrich B, Mosheimer B, Theurl M, Niederegger H, et al. Visfatin, an adipocytokine with proinflammatory and immunomodulating properties. *J Immunol* 2007;178:1748–1758.
- 103 Lee WJ, Wu CS, Lin H, Lee IT, Wu CM, Tseng JJ, et al. Visfatin-induced expression of inflammatory mediators in human endothelial cells through the NF- $\kappa$ B pathway. *Int J Obes* 2009;33:465–472.
- 104 Moschen AR, Kaser A, Enrich B, Mosheimer B, Theurl M, Niederegger H, et al. Visfatin, an adipocytokine with proinflammatory and immunomodulating properties. *J Immunol*. 2007, 178: 1748–1758.
- 105 Dahl TB, Yndestad A, Skjelland M, Oie E, Dahl A, Michelsen A, et al. Increased expression of visfatin in macrophages of human unstable carotid and coronary atherosclerosis: possible role in inflammation and plaque destabilization. *Circulation* 2007;115:972–980.
- 106 Brentano F, Schorr O, Ospelt C, Stanczyk J, Gay RE, Gay S, et al. Pre-B cell colony-enhancing factor/visfatin, a new marker of inflammation in rheumatoid arthritis with proinflammatory and matrix-degrading activities. *Arthritis Rheum* 2007;56:2829–2839.
- 107 Rho YH, Solus J, Sokka T, Oeser A, Chung CP, Gebretsadik T et al. Adipocytokines are associated with radiographic joint damage in rheumatoid arthritis. *Arthritis Rheum*. 2009, 60: 1906–1914
- 108 Gross JA, Johnston J, Mudri S, Enselman R, Dillon SR, Madden K et al. TACI and BCMA are receptors for a TNF homologue implicated in B-cell autoimmune disease. *Nature* 2000;404:995–9
- 109 Thompson JS, Bixler AS, Qian S, Vora K, Scott ML, Cachero TG et al. BAFF-R, a newly identified TNF receptor that specifically interacts with BAFF. *Science* 2001;293:2108–11
- 110 Yu-Hi K.; Choi BH, Cheon HG, DO MS: B cell activation factor (BAFF) is a novel adipokine that links obesity and inflammation. *Exp.Mol.Med* 2009, 41: 208–216
- 111 Zabel BA, Silverio AM., Butcher EC. Chemokine-like receptor 1 expression and chemerin- directed chemotaxis distinguish plasmacytoid from myeloid dendritic cells in human blood. *J.Immunol.*, 2005, 174: 244–251.
- 112 Hart R; Greaves DR. Chemerin contributes to inflammation by promoting adrophage adhesion to VCAM-1 and Fibronectin through clustering of VLA-4 and VLA-5. *J.Immunol*, 2010,185: 3728–3739
- 113 Berg V, Sveinbjörnsson B, Bendiksen S, Brox J, Meknas K, Figsenschau Y. Human articular chondrocytes express ChemR23 and chemerin; ChemR23 promotes inflammatory signalling upon binding the ligand chemerin<sup>21–157</sup>. *Arthr. Res & Ther* 2010, 12:R228
- 114 Iannone F; Lapadula G. Chemerin/ChemR23 pathway: a system beyond chemokines. *Arthr.Res.&Ther*. 2011, 13:104
- 115 Hart R, Greaves DR: Chemerin contributes to inflammation by promoting macrophage adhesion to VCAM-1 and fibronectin through clustering of VLA-4 and VLA-5. *J.Immunol*. 2010, 185: 3728–39
- 116 Yoshimura T; Oppenheim JJ. Chemerin reveals its chimeric nature. *J. Exp. Med*. 2008, 205 2187–90
- 117 Herenius M, Lebre C, Oliveira A, Wijbrandts C, Gerlag D, Tak PP: Chemerin levels are increased in rheumatoid arthritis patients who smoke, correlate with disease activity and decrease after Adalimumab. *Arthritis Rheum*. 2010, 63 (suppl.10), 1507
- 118 Nishida K, Okada Y, Nawata M, Saito K, Tanaka Y. Induction of hyperadiponectinemia following long-term treatment of patients with rheumatoid arthritis with infliximab (IFX), an anti-TNF- $\alpha$  antibody. *Endocr J* 2008;55:213–6.
- 119 Serelis J, Kontogianni MD, Katsiogiannis S, Bletsas M, Tektonidou MG, Skopouli FN. Effect of anti-TNF treatment on body composition and serum adiponectin levels of women with rheumatoid arthritis. *Clin Rheumatol* 2008;27:795–7.
- 120 Komai N, Morita Y, Sakuta T, Kuwabara A, Kashihara N. Anti-tumor necrosis factor therapy increases serum adiponectin levels with the improvement of endothelial dysfunction in patients with rheumatoid arthritis. *Mod Rheumatol* 2007;17:385–90.
- 121 Lewicki M, Kotyla P, Kucharz E. Increased adiponectin levels in women with rheumatoid arthritis after etanercept treatment. *J Rheumatol* 2009;36:1346–7.
- 122 Gonzalez-Gay MA, Garcia-Unzueta MT, Berja A, Gonzalez-Juanatey C, Miranda-Fillooy JA, Vazquez-Rodriguez TR, et al. Anti-TNF- $\alpha$

therapy does not modulate leptin in patients with severe rheumatoid arthritis. *Clin Exp Rheumatol* 2009;27:222–8.

- 123 Gonzalez-Gay MA, Vazquez-Rodriguez TR, Garcia-Unzueta MT, Berja A, Miranda-Fillooy JA, de Matias JM, et al. Visfatin is not associated with inflammation or metabolic syndrome in patients with severe rheumatoid arthritis undergoing anti-TNF-alpha therapy. *Clin Exp Rheumatol* 2010;28:56–62.
- 124 Gonzalez-Gay MA, Garcia-Unzueta MT, Berja A, Vazquez-Rodriguez TR, Gonzalez-Juanatey C, de Matias JM et al. Anti-TNF-alpha therapy modulates resistin in patients with rheumatoid arthritis. *Clin Exp Rheumatol* 2008;26:311–316.