

Hair-thread strangulation syndrome in childhood: a systematic review

Daniela Djokic^a, Gregorio P. Milani^{bc}, Sebastiano A. G. Lava^{de}, Gianluca Gualco^{af}, Teresa Corigliano^{af}, Mario G. Bianchetti^{af}, Camilla Lavagno^g

^a Pediatric Institute of Southern Switzerland, Ente Ospedaliero Cantonale, Bellinzona, Switzerland

^b Pediatric Unit, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy

^c Department of Clinical Sciences and Community Health, Università degli Studi di Milano, Milan, Italy

^d Pediatric Cardiology Unit, Department of Pediatrics, Centre Hospitalier Universitaire Vaudois and University of Lausanne, Lausanne, Switzerland

^e Heart Failure and Transplantation, Department of Pediatric Cardiology, Great Ormond Street Hospital, London, United Kingdom

^f Faculty of Biomedical Sciences, Università della Svizzera Italiana, Lugano, Switzerland

^g Pediatric Emergency Department, University Children's Hospital Zurich, Zurich, Switzerland

Summary

INTRODUCTION: Hair-thread strangulation syndrome describes the constriction of a body part by a tightly wound hair or thread. This research aims to review the literature about this entity.

METHODS: A systematic review was performed to characterise hair-thread strangulation syndrome in subjects aged ≤16 years. This pre-registered review (PROSPERO ID: CRD42022363996) followed the PRISMA methodology.

RESULTS: Subjects with digital strangulation were significantly younger (median = 4.0 [interquartile range: 2.0–6.1] months; n = 143) than females with genital strangulation (9.0 [6.8–11] years; n = 36), males with genital strangulation (5.1 [1.9–8.0] years; n = 36), and subjects with non-digital and non-genital strangulation (24 [13–48] months; n = 11). Digital strangulation was followed by an amputation in five (3.5%) and a reconstructive surgical intervention in seven (4.9%) cases. Sequelae occurred in four (11%) cases after female genital strangulation: clitoris autoamputation (n = 2) and surgical removal of a necrotic labium minus (n = 2). Severe complications were observed in 14 (39%) cases with male genital strangulation: urethral fistula (n = 7), urethral transection (n = 2), and partial penile autoamputation (n = 5). A partial uvular autoamputation was observed in one case (9.0%) with non-digital and non-genital strangulation.

CONCLUSIONS: Early recognition and management are crucial to avoid sequelae or long-term care in hair-thread strangulation syndrome.

Introduction

Hair-thread strangulation syndrome describes the acute constriction of a body part, such as digits or external genitalia, by a tightly wound hair or thread in a young infant, a child, or an adolescent [1]. Narrow constriction reduces lymphatic and venous drainage, resulting in pain, swelling, oedema, and, if recognised late, ischaemia [1]. This condi-

tion was first observed in the 19th century in a 4-week-old English boy with a painful penis swelling induced by a circumferential hair proximal to the swelling [2].

Hair-thread strangulation syndrome is either not mentioned or only briefly discussed in currently available paediatric textbooks. Its symptoms and presentation may be misdiagnosed, leading to delayed treatment and potentially adverse outcomes. A comprehensive literature review on this condition could help raise awareness among healthcare providers, ultimately improving diagnostic accuracy and patient outcomes. Therefore, we conducted a systematic review on this paediatric syndrome.

Methods

Search strategy

This literature review did not require Institutional Review Board approval. It was pre-registered at the International Prospective Register of Systematic Reviews (PROSPERO ID: CRD42022363996) and followed the 2020 edition [3] of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines (PRISMA) according to the Joanna-Briggs Manual. The search was undertaken in the National Library of Medicine, Excerpta Medica, and Web of Science databases without restriction. The search strategy combined the following terms: “hair strangulation syndrome” OR “hair tourniquet syndrome” OR “hair-thread strangulation syndrome” OR “hair-thread tourniquet syndrome” OR “strangulation syndrome” OR “tourniquet syndrome”.

Articles available in Google Scholar or already known to the authors were also considered. The search was undertaken in February 2022 and repeated before submission (November 2022).

Following a first round of selection based on title and abstract, the full text of the selected articles was evaluated for eligibility.

Dr. Gregorio P Milani
Fondazione IRCCS Ca'
Granda
Ospedale Maggiore Poli-
clinico
IT-20122 Milan
milani.gregorip[at]
gmail.com

Eligibility criteria and data extraction

Eligible articles were original reports and letters published in peer-review journals describing individual subjects aged ≤ 16 years with hair-thread strangulation affecting the digits, the external female genitalia, the penis, or other appendages. Cases with rubber band syndrome around the wrist were excluded [4]. Cases published only as abstracts and case series not detailing information on individual cases were also excluded.

The extracted data included demographics, the anatomic location, the source of strangulation (hair, threads, or elastic bands), the possible deliberate strangulation (by a caregiver or self-inflicted), the necessity of a reconstructive surgical procedure, and the outcome (development of long-term sequelae).

Two investigators independently (and in duplicate) performed the literature search, the selection of reports retained for analysis, and the data extraction in an unblinded manner with the support of a senior investigator. Any disagreements were discussed. One investigator entered the data into a pilot-tested database, and another verified them for accuracy.

Analysis

Pairwise deletion was used for missing data. Discrete variables are presented as counts and compared using Fisher's exact test. Numerical variables are presented as medians and interquartile ranges or as box-and-whisker plots and compared using Kruskal-Wallis and Tukey's tests [5, 6]. Two-sided P -values of <0.05 were considered statistically significant. Data were analysed using GraphPad Prism (version 9.5.1).

Results

Search output

The literature search identified 231 potentially relevant articles (figure 1). After removing irrelevant articles, 168 full-text articles were reviewed for eligibility. One hundred fifty-two articles [2, 7–157] describing individual cases of hair-thread strangulation syndrome published between 1951 and 2022 in English ($n = 147$), German ($n = 2$), French ($n = 1$), Spanish ($n = 1$), and Dutch ($n = 1$) were retained for the final analysis.

Fifty-eight articles reported cases from America, 45 from Asia, 39 from Europe, six from Oceania, and four from Africa. The articles provided information on 226 cases: 143 cases of digital strangulation [2, 7–90], 36 of female genital strangulation [22, 36, 49, 86, 91–119], 36 of male genital strangulation [7, 12, 24, 120–147], and 11 cases of non-digital and non-genital strangulation [148–157]. Characteristics of the individual cases are provided in the supplementary material (supplementary tables 1–4).

Clinical data

General data

Subjects with digital hair-thread strangulation syndrome (females: 50%; males: 50%) were significantly ($p < 0.01$) younger (4.0 [2.0–6.1] months) than females with genital

strangulation (9.0 [6.8–11] years), males with genital strangulation (5.1 [1.9–8.0] years), and subjects (females: 58%; males: 42%) with non-digital and non-genital strangulation (24 [13–48] months; figure 2). Females with genital strangulation, males with genital strangulation, and subjects with non-digital and non-genital strangulation did not differ significantly in age.

Twenty-nine (13%) of the 226 cases were explicitly reported as not induced by hairs but by threads arising from footed pyjamas, mittens, socks, underwear, and a nylon or elastic band. Three (1.3%) further cases were caused by hairs and threads.

The prevalence of cases not associated only with hairs was significantly higher ($p = 0.0003$) for digital strangulation ($n = 29$; 20%) than for the other forms of strangulation syndrome ($n = 3$; 3.6%).


A non-accidental cause of strangulation was suspected in 21 cases [12, 17, 36, 132, 133, 143]. Voluntary penile constriction was induced to manage enuresis in three cases [132, 133, 143].

Digital strangulation syndrome

Patients with digital hair-thread strangulation syndrome presented with irritability, crying, digital swelling, discolouration, and pain. About four fifths of the 143 cases of digital strangulation syndrome were observed on the toes (table 1). The female-to-male ratio did not differ significantly ($p = 0.336$) between infants with toe compared to finger strangulation syndrome. Cases with toe strangulation were slightly but significantly older than those with finger strangulation by 1.5 months on average ($p = 0.034$).


A detailed description of the affected digits was available for 136 (95%) of the 143 cases. Finger strangulation al-

Figure 1: Digital, female genital, male genital, and non-digital non-genital hair-thread strangulation syndrome in childhood. PRISMA flowchart of the literature search.



ways affected only one finger. In contrast, toe strangulation affected at least two toes in about a third of cases. The little finger, the big toe, and the little toe were relatively rarely

Figure 2: Age at presentation in paediatric patients with the various forms of hair-thread strangulation syndrome. The results are presented as box and whisker plots. Please note that the vertical axis is on a logarithmic scale. The age at presentation was significantly lower in children with digital strangulation than in children with strangulation of genitalia ($p < 0.005$) or other appendages ($p < 0.01$).



affected. Digital strangulation was induced by threads in eight (38%) of 21 cases with finger strangulation and in 12 (10%) of 115 cases with toe strangulation ($p = 0.0037$).

An amputation was performed in five (3.5%) cases: left thumb ($n = 1$), left middle toe ($n = 1$), right big toe ($n = 1$), right middle toe ($n = 1$), and right little toe ($n = 1$) [12, 26, 30, 53, 88]. A reconstructive surgical intervention was necessary in seven (4.9%) cases [12, 28, 29, 41, 70, 74, 81]. Removal of hair thread either mechanically or by applying a depilatory agent was followed by a full recovery in the remaining 131 (92%) cases.

Female genital strangulation syndrome

Hair-thread strangulation syndrome of the female external genital was reported in 36 cases, who presented with genital pain (sometimes associated with wide-base gait or painful micturition), vulvar swelling, or both genital pain and vulvar swelling. The strangulation was localised as follows (two structures were concurrently affected in two cases): clitoris ($n = 19$), labium minus ($n = 16$; bilateral in one case), labium majus ($n = 2$), and mons pubis ($n = 1$).

Sequelae were observed in four (11%) cases: total clitoris autoamputation in a 14-year-old adolescent, partial clitoris autoamputation in a 5-year-old girl, and surgical removal of a necrotic labium minus in a 6-year-old and a 10-year-old girl [95, 97, 105, 119]. The remaining 32 cases recovered uneventfully.

Male genital strangulation syndrome

Hair-thread strangulation syndrome of the male external genital was reported in 36 cases. Young infants presented with irritability and crying, often associated with swelling; older children presented with pain and swelling.

Table 1: Fingers and toes with hair-thread strangulation syndrome in 136 paediatric cases.

		Total	Right	Left
Fingers				
Female:male ratio	0.6*			
Age	2.4** [2.0–3.1] months			
n		21	10	11
Thumb		7	3	4
Forefinger		3	1	2
Middle finger		6	4	2
Ring finger		5	2	3
Little finger		0	0	0
One finger		21	10	11
Two fingers		0	0	0
Three fingers		0	0	0
Toes				
Female:male ratio	1.1			
Age	4.0 [2.6–5.0] months			
n		115	67	48
Big toe		3	2	1
Long toe		28	19	9
Middle toe		82	46	36
Ring toe		45	24	21
Little toe		8	5	3
One toe		79	47	32
Two toes		21	11	10
Three toes		15	9	6

* $p = 0.343$ and ** $p = 0.034$ versus toe strangulation.

Severe complications were observed in 14 (39%) cases [121, 123, 127, 128, 130, 134–136, 141, 143, 145–147]: urethral fistula ($n = 7$), urethral transection ($n = 2$), and partial penile autoamputation ($n = 5$).

Non-digital and non-genital strangulation syndrome

A non-digital and non-genital strangulation syndrome was noted in 11 cases (female-to-male ratio = 1.3): uvula ($n = 5$), neck ($n = 3$), circumvallate tongue papilla ($n = 2$), and tooth ($n = 1$). A partial uvular autoamputation was observed in one case [157].

Discussion

This literature review addressed the entire spectrum of conditions that may be caused by hair-thread strangulation in paediatric patients. While it found that they mainly affected the digits of young infants, they could also occur in older children. It also demonstrated that severe sequelae or the need for long-term care are possible (especially in cases with digital or penile strangulation). Furthermore, strangulation is sometimes non-accidental (e.g. intentional application to manage bedwetting). Finally, some cases were not caused by hairs but by threads or an elastic band.

Infants with hair strangulation often presented with a history of acute excessive crying [158, 159]. While the cause of acute and unexplained infantile crying is often benign, a careful history and physical examination remain the cornerstone of the evaluation. However, many cases present with local symptoms and signs such as pain, swelling, or discolouration. Detecting the encircling hair (which might be more challenging with light-coloured hairs) or thread is critical for diagnosis.

Because of the risk of ischemic damage, rapid management of hair-thread strangulation syndrome is crucial [160, 161]. Our review did not specifically address this issue. After taking a history and performing an examination to make the diagnosis, the first management step is currently appropriate analgesia [162]. Chemical depilatory agents, which break hairs (or threads) and subsequently lead to a weakening of the fibres and eventual dissolution, are currently the first-line treatment choice because they are generally painless and noninvasive [160, 161]. Mechanical release should be performed if strangulation release is not achieved after two attempts with a depilatory agent. Cases with strangulation near mucous membranes or with signs of ischemia should be treated with primary mechanical release [160, 161].

In infants, strangulation was caused mainly by maternal hair related to increased hair loss after birth [163]. It is assumed that hair may fall unnoticed into the bath water during routine care or diaper changing.

Our review had both weaknesses and strengths. Its main weakness was the low number of identified cases. The second weakness was its inability to address potential risk factors, such as socioeconomic status or ethnicity. Its most relevant strength was that it covered the entire spectrum of conditions that may be caused by hair-thread strangulation in paediatric patients. For example, two recent reviews focused explicitly on the hair-thread strangulation syndrome of the female genital tract or both the female and the male genital tracts [118, 164].

Conclusions

Early recognition and management are crucial to avoid sequelae or long-term care in hair-thread strangulation syndrome. Children and infants with unexpected acute onset of crying or local symptoms and signs such as pain, swelling, or discolouration require rapid medical evaluation. Healthcare professionals should be aware of this condition, with a particular focus on digital strangulation in infants and genital strangulation in children.

Data sharing statement

The data underlying this article can be shared upon reasonable request to the corresponding author.

Acknowledgments

Dr. Lava is the current recipient of research grants from Fonds de perfectionnement, Centre Hospitalier Universitaire Vaudois, Lausanne, Switzerland; Fondation SICPA, Prilly, Switzerland; Fondazione Dr. Ettore Balli, Bellinzona, Switzerland, Fondazione per il bambino malato della Svizzera italiana, Bellinzona, Switzerland and Frieda Locher-Hofmann Stiftung, Zürich, Switzerland.

Author contributions: Drs. Milani, Bianchetti and Lavagno conceptualized and designed the review. Drs. Djokic, Milani, Lava and Lavagno acquired the data and performed the analysis. Drs. Djokic, Bianchetti and Lavagno drafted the initial manuscript. All authors reviewed and revised the manuscript and approved the final manuscript as submitted.

Financial disclosure

This research received no external funding.

Potential competing interests

All authors have completed and submitted the International Committee of Medical Journal Editors form for disclosure of potential conflicts of interest. No potential conflict of interest related to the content of this manuscript was disclosed.

References

- Barton DJ, Sloan GM, Nicther LS, Reinisch JF. Hair-thread tourniquet syndrome. Pediatrics. 1988 Dec;82(6):925–8. <http://dx.doi.org/10.1542/peds.82.6.925>. PubMed. 0031-4005
- Ligation of the penis. Lancet. 1832;18(453):136. [http://dx.doi.org/10.1016/S0140-6736\(02\)82355-8](http://dx.doi.org/10.1016/S0140-6736(02)82355-8). 0140-6736
- Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. J Clin Epidemiol. 2021 Jun;134:178–89. <http://dx.doi.org/10.1016/j.jclinepi.2021.03.001>. PubMed. 1878-5921
- Meier R, Haug L, Surke C, Mathys L, Vögelin E. Acquired constriction ring: a case of rubber band syndrome. Pediatr Emerg Care. 2019 Jun;35(6):e113–5. <http://dx.doi.org/10.1097/PEC.0000000000000992>. PubMed. 1535-1815
- Bahna SL, McLarty JW. Data presentation. Ann Allergy Asthma Immunol. 2009 Oct;103(4 Suppl 1):S15–21. [http://dx.doi.org/10.1016/s1081-1206\(10\)60816-2](http://dx.doi.org/10.1016/s1081-1206(10)60816-2). [http://dx.doi.org/10.1016/S1081-1206\(10\)60816-2](http://dx.doi.org/10.1016/S1081-1206(10)60816-2). PubMed. 1081-1206
- Gaddis GM, Gaddis ML. Introduction to biostatistics: Part 5, Statistical inference techniques for hypothesis testing with nonparametric data [Erratum in: Ann Emerg Med 1991;20(5). Ann Emerg Med. 1990 Sep;19(9):1054–9. [http://dx.doi.org/10.1016/s0196-0644\(05\)82571-5](http://dx.doi.org/10.1016/s0196-0644(05)82571-5). PubMed. 0196-0644
- Alpert JJ, Filler R, Glaser HH. Strangulation of an appendage by hair wrapping. N Engl J Med. 1965 Oct;273(16):866–7. <http://dx.doi.org/10.1056/NEJM196510142731608>. PubMed. 0028-4793
- Curran JP. Digital strangulation by hair wrapping. J Pediatr. 1966 Jul;69(1):137–8. [http://dx.doi.org/10.1016/s0022-3476\(66\)80374-8](http://dx.doi.org/10.1016/s0022-3476(66)80374-8). [http://dx.doi.org/10.1016/S0022-3476\(66\)80374-8](http://dx.doi.org/10.1016/S0022-3476(66)80374-8). PubMed. 0022-3476
- Chown B. Digital strangulation by hair wrapping. Can Med Assoc J. 1967 Dec;97(25):1548. PubMed. 0008-4409
- Hill RH, James ES. Digital strangulation by hair wrapping. Can Med Assoc J. 1967 Nov;97(21):1293. PubMed. 0008-4409

11. Hack M, Brish M. A “finger equivalent” of the “toe tourniquet syndrome”. Pediatrics. 1972 Aug;50(2):348–9. <http://dx.doi.org/10.1542/peds.50.2.348>. PubMed. 0031-4005
12. Kerry RL, Chapman DD. Strangulation of appendages by hair and thread. J Pediatr Surg. 1973 Feb;8(1):23–7. [http://dx.doi.org/10.1016/0022-3468\(73\)90289-3](http://dx.doi.org/10.1016/0022-3468(73)90289-3). PubMed. 0022-3468
13. Mack JW, Takamoto RM, Jones FR, Zick HR. Toe tourniquet syndrome. West J Med. 1976 Oct;125(4):335–6. PubMed. 0093-0415
14. Douglas DD. Dissolving hair wrapped around an infant’s digit. J Pediatr. 1977 Jul;91(1):162. [http://dx.doi.org/10.1016/0022-3476\(77\)80478-2](http://dx.doi.org/10.1016/0022-3476(77)80478-2). PubMed. 0022-3476
15. Miller PR, Levi JH. Hair strangulation. J Bone Joint Surg Am. 1977 Jan;59(1):132. <http://dx.doi.org/10.2106/00004623-197759010-00028>. PubMed. 0021-9355
16. Arato A, Szamosfalvi I. Strangulation of toe caused by mother’s hair. Lancet. 1978 Jun;1(8076):1257–8. [http://dx.doi.org/10.1016/S0140-6736\(78\)92488-1](http://dx.doi.org/10.1016/S0140-6736(78)92488-1). PubMed. 0140-6736
17. Narkewicz RM. Distal digital occlusion. Pediatrics. 1978 Jun;61(6):922–3. <http://dx.doi.org/10.1542/peds.61.6.922>. PubMed. 0031-4005
18. Mosely LH. Digital entanglement in the blanket string: case of the necrosing digit. N Z Med J. 1979 Aug;90(642):150–2. PubMed. 0028-8446
19. Bérard J. Les cheveux étrangleurs ou toes tourniquet syndrome [Strangler hair or the toe tourniquet syndrome]. Pediatrie. 1981;36(7):539–42. PubMed. 0031-4021
20. Manfredi SJ. Accidental appendage strangulation. Plast Reconstr Surg. 1981 Jan;67(1):60–2. <http://dx.doi.org/10.1097/00006534-198101000-00014>. PubMed. 0032-1052
21. Collins AG. Hair-Thread Tourniquet syndrome. Australas J Dermatol. 1990;31(2):117–8. <http://dx.doi.org/10.1111/j.1440-0960.1990.tb00669.x>. PubMed. 0004-8380
22. Beasley SW. Strangulation of the clitoris by cotton fluff—another variant of hair-thread tourniquet syndrome. Pediatr Surg Int. 1991;6(4-5):383. <http://dx.doi.org/10.1007/BF00178666>. 0179-0358
23. Liow RY, Budny P, Regan PJ. Hair thread tourniquet syndrome. J Accid Emerg Med. 1996 Mar;13(2):138–9. <http://dx.doi.org/10.1136/emj.13.2.138>. PubMed. 1351-0622
24. Vázquez Rueda F, Núñez Núñez R, Gómez Meleno P, Blesa Sánchez E. Síndrome del torniquete por cabellos o hilos en dedos y pene [The hair-thread tourniquet syndrome of the toes and penis]. An Esp Pediatr. 1996 Jan;44(1):17–20. PubMed. 0302-4342
25. Conners G. Index of suspicion. Case 2. Hair tourniquet syndrome. Pediatr Rev. 1997 Aug;18(8):283–5. <http://dx.doi.org/10.1542/pir.18-8-283>. PubMed. 0191-9601
26. Leferink VJ, Klaase JM. Het teentourniquetsyndroom [Toe tourniquet syndrome]. Ned Tijdschr Geneesk. 1997 Dec;141(51):2499–501. PubMed. 0028-2162
27. Debo HF, Kostelanetz A, Morrison G, Brown EG. Red, edematous toes in two infants. J Pediatr Health Care. 1999;13(6 Pt 1):303. [http://dx.doi.org/10.1016/s0891-5245\(99\)90025-1](http://dx.doi.org/10.1016/s0891-5245(99)90025-1). [http://dx.doi.org/10.1016/S0891-5245\(99\)90025-1](http://dx.doi.org/10.1016/S0891-5245(99)90025-1). PubMed. 0891-5245
28. Sudhan ST, Gupta S, Plutarco C. Toe-tourniquet syndrome—accidental or intentional? Eur J Pediatr. 2000 Nov;159(11):866. [http://dx.doi.org/10.1007/PL00008357](http://dx.doi.org/10.1007/pl00008357). PubMed. 0340-6199
29. Sunil TM. The Hair-Thread-Tourniquet Syndrome - report of an unusual presentation of this rare condition. Hand Surg. 2001 Dec;6(2):231–3. <http://dx.doi.org/10.1142/s0218810401000746>. <http://dx.doi.org/10.1142/S0218810401000746>. PubMed. 0218-8104
30. Wang M, Schott J, Tunnessen WW Jr. Picture of the month. Hair-thread tourniquet syndrome. Arch Pediatr Adolesc Med. 2001 Apr;155(4):515–6. <http://dx.doi.org/10.1001/archpedi.155.4.515>. PubMed. 1072-4710
31. Corazza M, Carlà E, Altieri E, Virgili A. What syndrome is this? Tourniquet syndrome. Pediatr Dermatol. 2002;19(6):555–6. <http://dx.doi.org/10.1046/j.1525-1470.2002.00233.x>. PubMed. 0736-8046
32. Harris EJ. Acute digital ischemia in infants: the hair-thread tourniquet syndrome—a report of two cases. J Foot Ankle Surg. 2002;41(2):112–6. [http://dx.doi.org/10.1016/s1067-2516\(02\)80035-7](http://dx.doi.org/10.1016/s1067-2516(02)80035-7). [http://dx.doi.org/10.1016/s1067-2516\(02\)80035-7](http://dx.doi.org/10.1016/s1067-2516(02)80035-7). PubMed. 1067-2516
33. Smith AM, Peckett W, Davies M. Acquired constriction ring syndrome. Foot Ankle Int. 2003 Aug;24(8):640–1. <http://dx.doi.org/10.1177/107110070302400812>. PubMed. 1071-1007
34. Strahlman RS. Toe tourniquet syndrome in association with maternal hair loss. Pediatrics. 2003 Mar;111(3):685–7. <http://dx.doi.org/10.1542/peds.111.3.685>. PubMed. 1098-4275
35. Karalezli K, Karakoç Y, İlter S, Karalezli N, İrgit K. Hair-thread tourniquet syndrome. A case report. Acta Orthop Belg. 2004 Apr;70(2):180–1. PubMed. 0001-6462
36. Klusmann A, Lenard HG. Tourniquet syndrome—accident or abuse? Eur J Pediatr. 2004 Aug;163(8):495–8. <http://dx.doi.org/10.1007/s00431-004-1466-1>. PubMed. 0340-6199
37. Mackey S, Hettiaratchy S, Dickinson J. Hair-tourniquet syndrome—multiple toes and bilaterality. Eur J Emerg Med. 2005 Aug;12(4):191–2. <http://dx.doi.org/10.1097/00063110-200508000-00009>. PubMed. 0969-9546
38. Gutiérrez-Gómez C, Hernández-Antínez BG. Toes hair thread tourniquet syndrome. Case report. Bol Méd Hosp Infant México. 2006;63(4):268.0539-6115
39. Kurup HV, Gnanapavan M, McSweeney L. Hair-tourniquet syndrome: unwind or incise? Emerg Med Australas. 2006 Aug;18(4):415. <http://dx.doi.org/10.1111/j.1742-6723.2006.00882.x>. PubMed. 1742-6731
40. Lohana P, Vashishta GN, Price N. Toe-tourniquet syndrome: a diagnostic dilemma! Ann R Coll Surg Engl. 2006 Jul;88(4):W6–8. <http://dx.doi.org/10.1308/147870806X95276>. PubMed. 1478-7083
41. Mat Saad AZ, Purcell EM, McCann JJ. Hair-thread tourniquet syndrome in an infant with bony erosion: a case report, literature review, and meta-analysis. Ann Plast Surg. 2006 Oct;57(4):447–52. <http://dx.doi.org/10.1097/01.sap.0000222571.98387.71>. PubMed. 0148-7043
42. O’Quinn JC, Friedman RL, Wilms KL. Acquired constriction ring syndrome. J Am Podiatr Med Assoc. 2006;96(3):253–5. <http://dx.doi.org/10.7547/0960253>. PubMed. 8750-7315
43. von Känel OE. [Tourniquet syndrome — case report] [Tourniquet syndrome - case report]. Handchir Mikrochir Plast Chir. 2006 Apr;38(2):119–21. <http://dx.doi.org/10.1055/s-2006-923783>. PubMed. 0722-1819
44. Anwar MU, Ahmad M. The toe tourniquet syndrome. J Coll Physicians Surg Pak. 2007 May;17(5):309–10. PubMed. 1022-386X
45. Haene RA, Loeffler M. Hair tourniquet syndrome in an infant. J Bone Joint Surg Br. 2007 Feb;89(2):244–5. <http://dx.doi.org/10.1302/0301-620X.89B2.17626>. PubMed. 0301-620X
46. Kumaravel S. Chronic nonhealing ulcer in a finger of a toddler—a rare presentation of the hair thread tourniquet syndrome. Int J Dermatol. 2008 Nov;47(11):1174–6. <http://dx.doi.org/10.1111/j.1365-4632.2008.03871.x>. PubMed. 1365-4632
47. Singh V, Singh P, Sharma A, Sarkar J. Acquired constriction ring syndrome as a cause of inconsolable cry in a child: a case report. Cases J. 2008 Aug;1(1):92. <http://dx.doi.org/10.1186/1757-1626-1-92>. PubMed. 1757-1626
48. Al-Buainain H, Al-Jehani Y, Al-Ghamdi Z. Hair-thread tourniquet syndrome. Kuwait Med J. 2009;41(3):259–60.1607-8047
49. Cutrone M, Magagnato L. Hair thread tourniquet syndrome. Three cases in ten years. Eur J Pediatr Dermatol. 2009;19(1):23–6.1122-7672
50. García-Mata S, Hidalgo-Ovejero A. Hair tourniquet syndrome of the toe: report of 2 new cases. J Pediatr Orthop. 2009 Dec;29(8):860–4. <http://dx.doi.org/10.1097/BPO.0b013e3181b7ff14>. PubMed. 1539-2570
51. Ünlü C, Gundlach U, Vrouenraets BC. Toe hair-thread tourniquet syndrome. South Med J. 2009 Jun;102(6):663–4. <http://dx.doi.org/10.1097/SMJ.0b013e3181a2e33e>. PubMed. 1541-8243
52. Cevik Y, Kavalci C. Hair tourniquet syndrome. Ann Saudi Med. 2010;30(5):416–7. <http://dx.doi.org/10.4103/0256-4947.67088>. PubMed. 0975-4466
53. Heather J, Bisson M. Constriction “Band-Aid” syndrome causing digital ischaemia. N Z Med J. 2011 Jul;124(1338):107–10. PubMed. 1175-8716
54. O’Gorman A, Ratnapalan S. Hair tourniquet management. Pediatr Emerg Care. 2011 Mar;27(3):203–4. <http://dx.doi.org/10.1097/PEC.0b013e31820d65b7>. PubMed. 1535-1815
55. Onyeama CO, Vitale K, Cochran K, Onyeama GL. Swelling and redness of the fourth toe in a 3-month-old infant [Erratum in: Pediatr Rev. 2011;32(7)]. Pediatr Rev. 2011 Jun;32(6):253–5. <http://dx.doi.org/10.1542/pir.32-6-253>. <http://dx.doi.org/10.1542/pir.32-6-253>. PubMed. 0191-9601
56. Shankar R, Rao PS, Shetty KB, Sahu P, Singal R, Gupta S. Digital strangulation in a child: tourniquet syndrome. Indian J Dermatol Venereol Leprol. 2012;78(3):408. <http://dx.doi.org/10.4103/0378-6323.95489>. PubMed. 0973-3922
57. Sivathanan N, Vijayarajan L. Hair-thread tourniquet syndrome: a case report and literature review. Case Rep Med. 2012;2012:171368. <http://dx.doi.org/10.1155/2012/171368>. PubMed. 1687-9635
58. Wan WC, Yau MM, Lau DC, Huen KF, Chan LT. Hair tourniquet syndrome in a young infant. Hong Kong J Paediatr. 2012;17:119–22.2309-5393
59. Alvarez-Pérez A, Mateo S, Fernández-Redondo V, Toribio J. Hair-thread tourniquet syndrome: a hidden hazard. Pediatr Dermatol.

- 2013;30(4):e61–2. <http://dx.doi.org/10.1111/j.1525-1470.2012.01820.x>. PubMed. 1525-1470
60. Hickey BA, Gulati S, Maripuri SN. Hair toe tourniquet syndrome in a four-year-old boy. *J Emerg Med.* 2013 Feb;44(2):358–9. <http://dx.doi.org/10.1016/j.jemermed.2012.03.004>. PubMed. 0736-4679
61. Au S, Landers JT. Hair-thread tourniquet syndrome: a case report and review of treatment. *Dermatol Online J.* 2014;20(10):13030/qt7w85g4s4. <http://dx.doi.org/10.5070/D32010024259>.
62. Bizzotto N, Sandri A, Regis D, Carpeggiani G, Lavini F, Magnan B. Severe delayed hair toe tourniquet syndrome with bone erosion and flexor tendon lesion. *Cas Rep Pediatr.* 2014;2014:592323. <http://dx.doi.org/10.1155/2014/592323>. PubMed. 2090-6803
63. Kamal NM, Khan UU, Mirza SJ, Al-Malki TA. Toe tourniquet syndrome. *Saudi Med J.* 2014 Aug;35(8):865–7. PubMed. 1658-3175
64. Alruwaili N, Alshehri HA, Halimeh B. Hair tourniquet syndrome: successful management with a painless technique. *Int J Pediatr Adolesc Med.* 2015 Mar;2(1):34–7. <http://dx.doi.org/10.1016/j.ij-pam.2015.02.003>. PubMed. 2352-6467
65. Baloch N, Atif M, Rashid RH, Hashmi PM. Toe-tourniquet syndrome: A rare potentially devastating entity. *Malays Orthop J.* 2015 Nov;9(3):55–7. <http://dx.doi.org/10.5704/MOJ.1511.008>. PubMed. 1985-2533
66. Baştug O, Korkut S, Halis H, Kurtoğlu S. Hair-thread tourniquet syndrome in a preterm baby. *Turk Pediatri Ars.* 2015 Dec;50(4):245–7. <http://dx.doi.org/10.5152/TurkPediatri-Ars.2015.1301>. PubMed. 1306-0015
67. El-Kehdy J, Nasr J, Bitar C, El-Habr C, Abi-Rached Megarbane H, Sammour R, et al. Don't miss this blue toe: hair tourniquet syndrome. *J Eur Acad Dermatol Venereol.* 2015 Jan;29(1):191–2. <http://dx.doi.org/10.1111/jdv.12393>. PubMed. 1468-3083
68. Hussin P, Mawardi M, Masran MS, Ganaisan P. Hair tourniquet syndrome: revisited. *G Chir.* 2015;36(5):219–21. <http://dx.doi.org/10.11138/gchir/2015.36.5.219>. PubMed. 0391-9005
69. Kuiper JW, de Korte N. Hair thread tourniquet syndrome in a toe of an 18 mo old girl. *World J Clin Cases.* 2015 Apr;3(4):368–70. <http://dx.doi.org/10.12998/wjcc.v3.i4.368>. PubMed. 2307-8960
70. Pinkowsky GJ, Hennrikus WL. Toe tourniquet syndrome caused by cable tie. *Pediatr Emerg Care.* 2015 Jun;31(6):431–2. <http://dx.doi.org/10.1097/PEC.0000000000000461>. PubMed. 1535-1815
71. Pye S, Banks J, Pal C. Loom bands: a case of post-tourniquet syndrome. *Arch Dis Child.* 2015 May;100(5):481. <http://dx.doi.org/10.1136/archdischild-2014-307438>. PubMed. 1468-2044
72. van Meurs A, Korpershoek M, van Herwaarden-Lindenboom M. Female infant with toe swelling. *Ann Emerg Med.* 2016 Jun;67(6):785–96. <http://dx.doi.org/10.1016/j.annemergmed.2015.12.002>. PubMed. 1097-6760
73. Dunphy L, Verma Y, Morhij R, Lamyman M. Hair thread tourniquet syndrome in a male infant: a rare surgical emergency. *BMJ Case Rep.* 2017 Nov;2017:bcr2017221002. <http://dx.doi.org/10.1136/bcr-2017-221002>. PubMed. 1757-790X
74. Ojha S, Pandey VK, Garg G, Bansal R. "Missed band under the band": don't miss it! *Indian J Pediatr.* 2017 May;84(5):410–1. <http://dx.doi.org/10.1007/s12098-016-2275-z>. PubMed. 0973-7693
75. Rahul SK, Behera S, Suman P, Mohapatra S. Hair tourniquet syndrome. *Med J DY Patil Univ.* 2017;10(1):79–81. <http://dx.doi.org/10.4103/0975-2870.197919>. 0975-2870
76. Tengattini V, Magnano M, Patrizi A, Neri I. Sudden swelling and redness of the toe. *Clin Case Rep.* 2017 Sep;5(11):1901–2. <http://dx.doi.org/10.1002/ccr3.1115>. PubMed. 2050-0904
77. Uygur E, Çarkçı E, Ünkar E. Can washing socks without flipping inside out cause hair tourniquet syndrome? A claim with two case reports. *J Pediatr Orthop B.* 2017 Mar;26(2):193–4. <http://dx.doi.org/10.1097/BPB.0000000000000348>. PubMed. 1473-5865
78. Yadav R, Rahul SK, Kumar V, Zaheer H, Thakur K, Agarwal A. Toe tourniquet syndrome. *J Indira Gandhi Inst Med Sci.* 2017;3(1):34–5.
79. Booth J, Morse T. Hair toe tourniquets: a review of two case studies. *Emerg Nurse. Pediatr Int (Roma).* 2018;60(12):1095–6. <http://dx.doi.org/10.1111/ped.13712>. 0476-0069
80. Kesu Belani L, Leong JF, Narin Singh PSG, Abdullah S. Hair thread tourniquet syndrome in an infant: emergency exploration saves limbs. *2018;26(2):31-35.* <https://dx.doi.org/http://dx.doi.org/10.7748/en.2018.e1756>.
81. Sebaratnam DF, Hernández-Martín Á. Utility of ultrasonography in hair-thread tourniquet syndrome. *Pediatr Dermatol.* 2018 Mar;35(2):e138–9. <http://dx.doi.org/10.1111/pde.13400>. PubMed. 1525-1470
82. Singh AP, Tanger R, Mathur V, Gupta AK. Hair tourniquet syndrome: A rare case report. *Menoufia Med J.* 2018;30(4):1260–1. http://dx.doi.org/10.4103/mmj.mmj_111_17. 1110-2098
83. Aghbari FZ. Le syndrome des cheveux étrangleurs: à propos d'un cas [Strangler hair syndrome: about a case]. *Pan Afr Med J.* 2019 Jul;33:266. <http://dx.doi.org/10.11604/pamj.2019.33.266.15699>. PubMed. 1937-8688
84. Aslantürk O, Özbeyp R, Yılmaz Ö, Ergen E. Hair tourniquet syndrome of toes and fingers in infants. *Acta Orthop Traumatol Turc.* 2019 Jul;53(4):306–9. <http://dx.doi.org/10.1016/j.aott.2019.04.010>. PubMed. 2589-1294
85. Basheer SM, Gerashi AR, Basith MA. Hair tourniquet syndrome, an unusual presentation: A case report and review of literature. *Int J Case Rep Images.* 2019;10:100992Z01BS2019. <https://dx.doi.org/http://dx.doi.org/100992Z01BS2019>.
86. Ihara T, Takei H, Kishibe S, Nomura O. Hair tourniquet syndrome on the toe and labia. *Cureus.* 2019;11(12):e6377. <http://dx.doi.org/10.7759/cureus.6377>. PubMed. 2168-8184
87. Azizrahman A, Wani SA, Alnasser OM. Irritable infant: a rare and easy to miss case for strangulated toe. *Saudi J Emerg Med.* 2020;1(2):121–3. <http://dx.doi.org/10.24911/SJEMED.72-1573244443>. 1658-8487
88. McGrath RB, Cahill KC, Eadic PA, Harty SB. Thumb autoamputation following tourniquet syndrome. *BMJ Case Rep.* 2020 Mar;13(3):e233143. <http://dx.doi.org/10.1136/bcr-2019-233143>. PubMed. 1757-790X
89. Shafiq MA, Anwar K, Tan JA. The fretful child with swollen appendages: mitten the insidious. *Asian J Med Biomed.* 2020;4(1):31–3. <http://dx.doi.org/10.37231/ajmb.2020.4.1.327>.
90. Suteu M, Yıldırın G, Akdag O. Hair-thread tourniquet syndrome: two case reports. *Turk J Plast Surg.* 2020;28:237–9. http://dx.doi.org/0.4103/tjps.tjps_62_19.
91. Chapman HL. Digital strangulation by hair wrapping. *Can Med Assoc J.* 1968 Jan;98(2):125. PubMed. 0008-4409
92. Press S, Schachner L, Paul P. Clitoris tourniquet syndrome. *Pediatrics.* 1980 Nov;66(5):781–2. <http://dx.doi.org/10.1542/peds.66.5.781>. PubMed. 0031-4005
93. Summers JL, Guira AC. Hair strangulation of the external genitalia: report of two cases. *Ohio State Med J.* 1973 Sep;69(9):672–3. PubMed. 0030-1124
94. Webley JA, Schleif DR, Coleman J. Tourniquet Syndrome: an unusual presentation. *Ann Emerg Med.* 1981 Sep;10(9):494–5. [http://dx.doi.org/10.1016/s0196-0644\(81\)80288-0](http://dx.doi.org/10.1016/s0196-0644(81)80288-0). [http://dx.doi.org/10.1016/S0196-0644\(81\)80288-0](http://dx.doi.org/10.1016/S0196-0644(81)80288-0). PubMed. 0196-0644
95. Rich MA, Keating MA. Hair tourniquet syndrome of the clitoris. *J Urol.* 1999 Jul;162(1):190–1. <http://dx.doi.org/10.1097/00005392-199907000-00067>. PubMed. 0022-5347
96. Sylwestrzak MS, Fischer BF, Fischer H. Recurrent clitoral tourniquet syndrome. *Pediatrics.* 2000 Apr;105(4 Pt 1):866–7. <http://dx.doi.org/10.1542/peds.105.4.866>. PubMed. 0031-4005
97. Kuo JH, Smith LM, Berkowitz CD. A hair tourniquet resulting in strangulation and amputation of the clitoris. *Obstet Gynecol.* 2002 May;99(5 Pt 2):939–41. [http://dx.doi.org/10.1016/s0029-7844\(01\)01753-7](http://dx.doi.org/10.1016/s0029-7844(01)01753-7). [http://dx.doi.org/10.1016/S0029-7844\(01\)01753-7](http://dx.doi.org/10.1016/S0029-7844(01)01753-7). PubMed. 0029-7844
98. Serour F, Gorenstein A. Treatment of the toe tourniquet syndrome in infants. *Pediatr Surg Int.* 2003 Oct;19(8):598–600. <http://dx.doi.org/10.1007/s00383-003-1034-1>. PubMed. 0179-0358
99. Bacon JL, Burgis JT. Hair thread tourniquet syndrome in adolescents: a presentation and review of the literature. *J Pediatr Adolesc Gynecol.* 2005 Jun;18(3):155–6. <http://dx.doi.org/10.1016/j.jpag.2005.03.010>. PubMed. 1083-3188
100. Golshovsky J, Chuen J, Tung PH. Hair-thread tourniquet syndrome. *J Paediatr Child Health.* 2005 Mar;41(3):154–5. <http://dx.doi.org/10.1111/j.1440-1754.2005.00569.x>. PubMed. 1034-4810
101. Osterhoudt KC, Beleniski F. Images in emergency medicine. The hair tourniquet syndrome. *Ann Emerg Med.* 2005 Apr;45(4):458–63. <http://dx.doi.org/10.1016/j.annemergmed.2004.08.051>. PubMed. 1097-6760
102. Alverson B. A genital hair tourniquet in a 9-year-old girl. *Pediatr Emerg Care.* 2007 Mar;23(3):169–70. <http://dx.doi.org/10.1097/PEC.0b013e3180328cc4>. PubMed. 1535-1815
103. Pomeranz M, Schachter B, Capua T, Beyth Y. Hair-thread tourniquet syndrome of labia minor. *J Pediatr Adolesc Gynecol.* 2009 Oct;22(5):e111–3. <http://dx.doi.org/10.1016/j.jpag.2008.07.011>. PubMed. 1873-4332
104. Stock C, Wang LC, Spigland NA. Untangling a web: an unusual case of labial necrosis in an adolescent female. *J Pediatr Adolesc Gynecol.* 2012 Apr;25(2):e21–2. <http://dx.doi.org/10.1016/j.jpag.2011.11.001>. PubMed. 1873-4332

105. Bannier MA, Miedema CJ. Hair tourniquet syndrome. *Eur J Pediatr.* 2013 Feb;172(2):277. <http://dx.doi.org/10.1007/s00431-012-1883-5>. PubMed. 1432-1076
106. Dua A, Jamshidi R, Lal DR. Labial hair tourniquet: unusual complication of an unrepaired genital laceration. *Pediatr Emerg Care.* 2013 Jul;29(7):829-30. <http://dx.doi.org/10.1097/PEC.0b013e3182986184>. PubMed. 1535-1815
107. Lee JH, Pringle K, Rajimwale A. Red herring vaginal discharge. *BMJ Case Rep.* 2013 Sep;2013 sep18 1:bcr2013200367. <http://dx.doi.org/10.1136/bcr-2013-200367>. PubMed. 1757-790X
108. Ozcakir E, Kaya M. A genital hair tourniquet syndrome: case report and review of the literature. *Pediatr Urol Case Rep.* 2014;1(2):1-4. <http://dx.doi.org/10.14534/PUCR.201424403>. 2148-2969
109. Panagidis A, Sinopidis X, Alexopoulos V, Georgiou G. Hair tourniquet syndrome of labia minora. *APSP J Case Rep.* 2015 May;6(2):22. PubMed. 2218-8185
110. Parlak M, Karakaya AE. Hair-thread tourniquet syndrome of the hypertrophic clitoris in a 6-year-old girl. *Pediatr Emerg Care.* 2015 May;31(5):363-4. <http://dx.doi.org/10.1097/PEC.0000000000000341>. PubMed. 1535-1815
111. Zimmerman LN, Wagner AJ. Clitoral hair tourniquet: a case report and review of the literature. *Int J Pediatr Res.* 2015;1(2):007. <http://dx.doi.org/10.23937/2469-5769/1510007>.
112. Zhong N, Bhattacharya R. Hair tourniquet of the clitoris: A novel treatment. *J Pediatr Surg Case Rep.* 2017;27:26-7. <http://dx.doi.org/10.1016/j.epsc.2017.09.003>. 2213-5766
113. Roman E. Adolescent labial hair tourniquet. *Aust N Z J Obstet Gyneacol.* 2018;58:72.0004-8666
114. Erikci VS, Sayan A, Gokhan Koyluoglu G. Hair tourniquet syndrome of the clitoris and labia minora: report of three cases and review of the literature. *Pediatr Urol Case Rep.* 2019;6(6):127-31. <http://dx.doi.org/10.14534/j-pucr.2019655917>. 2148-2969
115. Imam AS, Florez CE, Beland L, Swana H, Rich M. Hair tourniquet of the labia minora. *J Pediatr Surg Case Rep.* 2019;44:101160. <http://dx.doi.org/10.1016/j.epsc.2019.01.012>. 2213-5766
116. Diaz-Morales O, Martinez-Pajares JD, Ramos-Diaz JC, Gutierrez Del Alamo Lopez J, Trigo-Moreno J. Genital hair-thread tourniquet syndrome. *J Pediatr Adolesc Gynecol.* 2020 Dec;33(6):715-9. <http://dx.doi.org/10.1016/j.jpag.2020.05.009>. PubMed. 1873-4332
117. Minodier P, Cohen-Boulakia R, Aguesse C, Fruscione S, Urbina D. Genital hair tourniquet syndrome in a 5-year old girl. *Ann Clin Med Case Reports.* 2020;5(5):1-2. <http://dx.doi.org/10.47829/ACM-CR.2020.5504>.
118. Adjei N, Lynn A, Dugan K, Vash-Margita A. Medically Indicated Labiaplasty in a Case of Recurrent Labial Hair Thread Tourniquet Syndrome in a Premenarchal Girl. *J Pediatr Adolesc Gynecol.* 2021 Oct;34(5):761-3. <http://dx.doi.org/10.1016/j.jpag.2021.03.007>. PubMed. 1873-4332
119. Chawla RK, Lakshminarayanan S, Jay MS. Vulvar discomfort in a 10-year-old girl. *Pediatr Rev.* 2021 Aug;42(8):e33-6. <http://dx.doi.org/10.1542/pir.2020-003020>. PubMed. 1526-3347
120. DeWEESE JA, Eikner WC. Strangulation of the penis with a human hair. *Urol Cutaneous Rev.* 1951 Jan;55(1):37-8. PubMed. 0096-6878
121. Farah R, Cerny JC. Penis tourniquet syndrome and penile amputation. *Urology.* 1973 Sep;2(3):310-1. [http://dx.doi.org/10.1016/0090-4295\(73\)90473-1](http://dx.doi.org/10.1016/0090-4295(73)90473-1). PubMed. 0090-4295
122. Haddad FS. Penile strangulation by human hair. Report of three cases and review of the literature. *Urol Int.* 1982;37(6):375-88. <http://dx.doi.org/10.1159/000280843>. PubMed. 0042-1138
123. McClure WJ, Gradinger GP. Hair strangulation of the glans penis. *Plast Reconstr Surg.* 1985 Jul;76(1):120-3. <http://dx.doi.org/10.1097/00006534-198507000-00022>. PubMed. 0032-1052
124. Sheinfeld J, Cos LR, Erturk E, Cockett AT. Penile tourniquet injury due to a coil of hair. *J Urol.* 1985 Jun;133(6):1042-3. [http://dx.doi.org/10.1016/s0022-5347\(17\)49371-2](http://dx.doi.org/10.1016/s0022-5347(17)49371-2). PubMed. 0022-5347
125. Mariani PJ, Wagner DK. Topical cocaine prior to treatment of penile tourniquet syndrome. *J Emerg Med.* 1986;4(3):205-7. [http://dx.doi.org/10.1016/0736-4679\(86\)90042-9](http://dx.doi.org/10.1016/0736-4679(86)90042-9). PubMed. 0736-4679
126. Pantuck AJ, Kraus SL, Barone JG. Hair strangulation injury of the penis. *Pediatr Emerg Care.* 1997 Dec;13(6):423-4. <http://dx.doi.org/10.1097/00006565-199712000-00020>. PubMed. 0749-5161
127. Bangroo AK, Chauhan S. Hair tourniquet syndrome. *J Indian Assoc Pediatr Surg.* 2005;10(1):55-6. <http://dx.doi.org/10.4103/0971-9261.16080>. 0971-9261
128. Chaware SM, Gajbhiye R, Singh RA. Penile tourniquet injury due to a coil of hair. *Indian J Plast Surg.* 2006;39(1):70-2. <http://dx.doi.org/10.1055/s-0039-1700466>. 0970-0358
129. Dar NR, Siddiqui S, Qayyum R, Ghafoor T. Hair coil strangulation—an uncommon cause of penile edema. *Pediatr Dermatol.* 2007;24(4):E33-5. <http://dx.doi.org/10.1111/j.1525-1470.2007.00436.x>. PubMed. 1525-1470
130. Hussain HM. A hair tourniquet resulting in strangulation and amputation of penis: case report and literature review. *J Paediatr Child Health.* 2008 Oct;44(10):606-7. <http://dx.doi.org/10.1111/j.1440-1754.2008.01394.x>. PubMed. 1034-4810
131. Okeke LI. Thread embedded into penile tissue over time as an unusual hair thread tourniquet injury to the penis: a case report. *J Med Case Rep.* 2008 Jul;2(1):230. <http://dx.doi.org/10.1186/1752-1947-2-230>. PubMed. 1752-1947
132. Basner A, Gumpert S, Lütkenhaus D, Kaatsch HJ, Schwark T. Penishämatom mit Schwellung. Seltener Fall von Kindesmisshandlung? Rechtsmedizin [Penile hematoma with swelling. A rare case of child abuse?]. 2012;22:479-481. <https://dx.doi.org/https://dx.doi.org/10.1007/s00194-012-0852-7>. <http://dx.doi.org/10.1007/s00194-012-0852-7>
133. Pahwa HS, Kumar A, Srivastava R, Kumar S, Goel A, Ahmad A. Partial penile amputation due to penile tourniquet syndrome in a child troubled with primary nocturnal enuresis—a rare emergency. *Urology.* 2013 Mar;81(3):653-4. <http://dx.doi.org/10.1016/j.urology.2012.11.012>. PubMed. 1527-9995
134. Jesus LE, Braga JJ, Rocha JM, Dekermacher S, Anderson KM. Penile hair tourniquet resulting in hypospadias failure. *Indian J Urol.* 2014 Jul;30(3):345-7. <http://dx.doi.org/10.4103/0970-1591.128508>. PubMed. 0970-1591
135. Dikshit VK, Gupta RK, Kothari PR, Gupta AR, Kamble RS, Kesan KV. Near total penile amputation caused by hair tourniquet, managed with the URAGPI procedure. *Afr J Urol.* 2015;21(4):254-7. <http://dx.doi.org/10.1016/j.afju.2014.11.010>. 1110-5704
136. Gazali ZA, Singal AK. Hair Thread Tourniquet Syndrome of Penis Causing Urethral Fistula. *Indian Pediatr.* 2015 Jun;52(6):538. PubMed. 0974-7559
137. Zengin K, Ozdamar MY, Albayrak S, Tanik S, Atar M, Bakirtas H, et al. Hair coil penile tourniquet syndrome in an unusual age. *Case Rep Urol.* 2015;2015:642547. <http://dx.doi.org/10.1155/2015/642547>. PubMed. 2090-696X
138. Sekhon V, Peters NJ, Mahajan JK. Penile hair tourniquet syndrome: A case report and review of literature. *Pediatr Urol Case Rep.* 2017;4(3):311-4. <http://dx.doi.org/10.14534/PUCR.2017327011>. 2148-2969
139. Sheikh Andalibi MS, Azfarfar A, Esmaeeli M, Sheyk Andalibi N, Alizadeh A, Ravanshad Y. Penile hair tourniquet syndrome due to a coil of hair: first report in Iran. *Nephrourol Mon.* 2017;9(4):e59788. <http://dx.doi.org/10.5812/numphonthly.59788>. 2251-7006
140. Özçift B, Ağras K. Hair tourniquet syndrome of penis: A rare situation in boys with serious complications if not recognized. *Turk J Urol.* 2018 Aug;45(4):322-4. <http://dx.doi.org/10.5152/tud.2018.36699>. PubMed. 2149-3235
141. Arshadi H, Fatemi Behbehani SM, Karimi A, Lotfi B. Penile strangulation by hair: a case report. *Case Rep Clin Pract.* 2019;4(2):44-7. <http://dx.doi.org/10.18502/crcp.v4i2.1705>. 2538-2691
142. Degheili JA, El-Asmar JM, Aoun B. Penile hair strangulation: A prodigious cause of pediatric agitation. *Clin Case Rep.* 2019 May;7(7):1452-3. <http://dx.doi.org/10.1002/ccr3.2198>. PubMed. 2050-0904
143. Bouassida K, Ben Ahmed K, Ben Othmen M, Jaidane M. Penile hair coil strangulation in a 9-year-old patient: surgical management and review of the literature. *Ann Med Surg (Lond).* 2020 Oct;60:50-5. <http://dx.doi.org/10.1016/j.amsu.2020.10.020>. PubMed. 2049-0801
144. Rawls WF, White JT, Mohamed A, Peppas D, Rosenberg E. Case report: penile strangulation secondary to hair tourniquet. *Front Pediatr.* 2020 Aug;8:477. <http://dx.doi.org/10.3389/fped.2020.00477>. PubMed. 2296-2360
145. Patterson JW, Haberman C, Skelton JA, Pranikoff T, Northrop S. Infantile penile degloving injury secondary to hair tourniquet. *J Pediatr Surg Case Rep.* 2022;80:102250. <http://dx.doi.org/10.1016/j.epsc.2022.102250>. 2213-5766
146. Rampersad B, Dubay C, Doodnath R. Penile hair tourniquet to base of penis resulting in circumferential separation of shaft skin without vascular compromise: an unusual case report. *Urol Case Rep.* 2022 Feb;42:102030. <http://dx.doi.org/10.1016/j.eucr.2022.102030>. PubMed. 2214-4420
147. Loloj I, Andreas D, Gitlin J. Severe penile hair tourniquet syndrome with near amputation of the glans. *J Clin Urol.* 2023 Jan;16(1):51-4. <http://dx.doi.org/10.1177/205145820939465>. ; Epub ahead of print. 2051-4158

148. McNeal RM, Cruickshank JC. Strangulation of the uvula by hair wrapping. Clin Pediatr (Phila). 1987 Nov;26(11):599–600. <http://dx.doi.org/10.1177/000992288702601111>. PubMed. 0009-9228
149. Krishna S, Paul RJ. Hair tourniquet of the uvula. J Emerg Med. 2003 Apr;24(3):325–6. [http://dx.doi.org/10.1016/s0736-4679\(02\)00760-6](http://dx.doi.org/10.1016/s0736-4679(02)00760-6). PubMed. 0736-4679
150. Chegwidden HJ, Poirier MP. Near strangulation as a result of hair tourniquet syndrome. Clin Pediatr (Phila). 2005 May;44(4):359–61. <http://dx.doi.org/10.1177/000992280504400413>. PubMed. 0009-9228
151. Schneider K, Kennebeck S, Madden L, Campbell A. Hair tourniquet of the circumvallate papillae: a potentially “hairy” situation. Pediatr Emerg Care. 2013 Aug;29(8):924–5. <http://dx.doi.org/10.1097/PEC.0b013e31829ec4c7>. PubMed. 1535-1815
152. Flores JR. Hair tourniquet syndrome in the dental patient. Anesth Prog. 2014;61(3):111–2. <http://dx.doi.org/10.2344/0003-3006-61.3.111>. PubMed. 0003-3006
153. Alghamdi F, Alrashed R, Alsadhan N. Tongue papillae thread tourniquet removal under sedation: an apparently challenging case. J Gen Emerg Med. 2018;3(2):039.
154. Wong K, Tiyagura G, Langhan ML. Accidental strangulation: a rare but potential risk of co-sleeping. Pediatr Oncall. 2019;16(1):21–2. <http://dx.doi.org/10.7199/ped.oncall.2019.2>. 0973-0966
155. Stansfield J, Coey J, Mirza O, Loughran S. Uvula strangulation: a rare case of hair thread tourniquet syndrome. Int J Surg Case Rep. 2020;2(3):2. <http://dx.doi.org/10.31487/j.IJSCR.2020.03.10>. 2674-4171
156. Chan LY, Chan TW, Tay MK. Accidental hair strangulation of a toddler caught on home surveillance camera. Arch Dis Child. 2021 Nov;106(11):1046. <http://dx.doi.org/10.1136/archdischild-2020-320669>. PubMed. 1468-2044
157. Leighton J, Powell J, McLarnon C. Hair-thread Tourniquet Syndrome causing uvular strangulation in a child. BMJ Case Rep.
- 2021 Apr;14(4):e242350. <http://dx.doi.org/10.1136/bcr-2021-242350>. PubMed. 1757-790X
158. Poole SR. The infant with acute, unexplained, excessive crying. Pediatrics. 1991 Sep;88(3):450–5. <http://dx.doi.org/10.1542/peds.88.3.450>. PubMed. 0031-4005
159. Freedman SB, Al-Harthi N, Thull-Freedman J. The crying infant: diagnostic testing and frequency of serious underlying disease. Pediatrics. 2009 Mar;123(3):841–8. <http://dx.doi.org/10.1542/peds.2008-0113>. PubMed. 1098-4275
160. Gottlieb M, Holladay D, Spearman D. Current approach to the evaluation and management of hair-thread tourniquets. Pediatr Emerg Care. 2019 May;35(5):377–9. <http://dx.doi.org/10.1097/PEC.0000000000001827>. PubMed. 1535-1815
161. Kudzinskas A, Reed A, Mahdi M, Tyler M. Evaluation of depilatory agents in the treatment of hair-thread tourniquet syndrome. J Emerg Med. 2021 Nov;61(5):507–16. <http://dx.doi.org/10.1016/j.jemermed.2021.07.013>. PubMed. 0733-8627
162. Bauman BH, McManus JG Jr. Pediatric pain management in the emergency department. Emerg Med Clin North Am. 2005 May;23(2):393–414. <http://dx.doi.org/10.1016/j.emc.2004.12.008>. PubMed. 1060-0280
163. Eastham JH. Postpartum alopecia. Ann Pharmacother. 2001 Feb;35(2):255–8. <http://dx.doi.org/10.1345/aph.10153>. PubMed. 1060-0280
164. De Vitis LA, Barba M, Lazzarin S, Molinari S, Spinelli M, Arosio E, et al. Female genital hair-thread tourniquet syndrome: a case report and literature systematic review. J Pediatr Adolesc Gynecol. 2021 Feb;34(1):65–70. <http://dx.doi.org/10.1016/j.jpag.2020.07.007>. PubMed. 1873-4332

Appendix: Supplementary tables

Supplementary table 1:

Characteristics of 143 individuals affected by digital hair-thread strangulation syndrome. The country of publication is identified using the International vehicle registration code.

Case	First author	Year of publication	Country	Sex	Age (months)
1	Alpert JJ [7]	1965	USA	Male	2.0
2				Male	19
3	Curran JP [8]	1966	USA	Male	2.5
4	Chonwn [9]	1967	CDN		12
5	Hill RH [10]	1967	CDN		5.0
6	Hack M [11]	1972	IL	Female	0.8
7	Kerry RL [12]	1973	USA	Male	4.0
8				Male	1.8
9				Male	5.0
10				Female	2.3
11				Male	1.5
12				Female	1.8
13				Female	3.5
14	Mack JW [13]	1976	USA	Female	11
15				Male	5.0
16	Douglas DD [14]	1977	USA		10
17				Female	11
18	Miller PR [15]	1977	USA	Female	15
19	Arato A [16]	1978	H		6.0
20	Narkewicz RM [17]	1978	USA	Male	4.0
21				Male	3.0
22				Male	1.0
23	Mosely LH [18]	1979	NZ	Female	2.0
24	Bérard J [19]	1981	F	Male	6.0
25				Male	3.0
26				Male	3.0
27	Manfemaleredi SJ [20]	1981	USA	Female	7.0
28				Male	3.0
29	Barton DJ [1]	1988	USA	Male	5.0
30				Female	2.0
31				Female	4.0
32				Male	1.8
33				Male	144
34				Female	0.6
35	Collins AG [21]	1990	AUS	Female	14
36	Beasley SW [22]	1993	USA		2.5
37	Liow RY [23]	1996	UK	Female	18
38				Male	3.0
39	Vazquez Rueda F [24]	1996	E	Female	2.0
40				Male	4.0
41				Male	2.0
42				Female	5.0
43				Male	3.0
44				Male	7.0
45	Connors G [25]	1997	USA	Male	5.0
46	Lefemaleerink VJ [26]	1997	NL	Male	4.0
47				Male	4.0
48				Female	2.0
49				Male	0.8
50	Debo HF [27]	1999	USA	Male	2.0
51				Male	4.0
52	Sudhan ST [28]	2000	UK		5.0
53	Sunil TM [29]	2001	IND	Female	8.0
54	Wang M [30]	2001	USA		1.5
55	Corazza M [31]	2002	I	Female	72
56	Harris EJ [32]	2002		Female	5.0
57				Male	5.0
58	Smith AM [33]	2003	UK	Male	7.0
59	Strahlman RS [34]	2003	USA	Male	3.0
60	Karalezli K [35]	2004	B	Male	3.0
61	Klusmann A [36]	2004	D		4.0

62					4.0
63					1.5
64					2.0
65	Mackey S [37]	2005	UK	Female	3.0
66	Gutiérrez-Gómez C [38]	2006	MEX	Female	2.0
67	Kurup HV [39]	2006	UK	Male	3.0
68	Lohana P [40]	2006	UK		1.5
69	Mat Saad AZ [41]	2006	IRL	Female	3.0
70	O'Quinn JC [42]	2006	USA	Female	2.3
71	von Känel OE [43]	2006	CH	Male	2.0
72	Anwar MU [44]	2007	UK	Female	4.0
73	Haene RA [45]	2007	UK		2.8
74	Kumaravel S [46]	2008	IND		24
75	Singh V [47]	2008	UK	Female	14
76	Al-Buainain H [48]	2009	KSA	Male	3.0
77	Cutrone M [49]	2009	I	Female	2.0
78				Female	2.0
79	Garcia-Mata S [50]	2009	USA	Female	18
80				Male	5.0
81	Ünlü C [51]	2009	NL	Male	5.0
82	Cevik Y [52]	2010	TR	Female	4.0
83	Heather J [53]	2011	NZ	Female	36
84	O'Gorman A [54]	2011	USA	Male	3.0
85	Onyeama CO [55]	2011	USA	Male	3.0
86	Sivathasan N [57]	2012	UK		
87					
88	Shankar R [56]	2012	IND	Female	1.0
89	Wan WCY [58]	2012	CHN	Female	2.0
90	Alvarez-Perez A [59]	2013	E	Male	8.0
91	Hickey BA [60]	2013	UK	Male	48
92	Au S [61]	2014	USA	Male	6.0
93	Bizzotto N [62]	2014	I	Female	2.0
94	Kamal NM [63]	2014	KSA	Female	6.0
95				Male	6.0
98	Baloch N [65]	2015	PK	Female	4.0
99	Bastug O [66]	2015	TR	Female	0.5
100	EI-Kehdy J [67]	2015	RL	Female	6.0
101					
102	Hussin P [68]	2015	MAL	Female	3.0
103				Male	5.0
104	Kuiper JW [69]	2015	NL	Female	18
105	Pinkowsky GJ [70]	2015	USA	Female	84
106	Pye S [71]	2015	UK	Female	48
	Alruwaili N [64]	2016	KSA	Male	6.0
					0.5
107	van Meurs A [72]	2016	USA	Female	4.0
108	Dunphy L [73]	2017	UK	Male	2.0
109	Ojha S [74]	2017	IND	Female	12
110	Rahul SK [75]	2017	IND	Male	3.0
111	Tengattini V [76]	2017	I	Female	8.0
112	Uygur E [77]	2017	TR	Female	4.0
113				Female	6.0
114	Yadav R [78]	2017	IND	Male	2.0
115	Booth J [79]	2018	UK	Male	3.0
116					14
117	Ihara T [86]	2018	J	Female	1.3
118	Sebaratnam D [81]	2018	E	Female	24
119	Singh AP [82]	2018	IND	Male	1.5
120	Agherbi F [83]	2019	MA		
121	Aslantürk O [84]	2019	TR	Male	2.0
122				Male	5.6
123				Female	1.9
124				Female	3.5
125				Female	0.6

126				Male	4.3
127				Male	2.0
128				Female	4.0
129				Male	1.0
130				Male	13
131				Male	0.8
132				Male	3.9
133				Female	6.0
134				Female	2.8
135				Female	2.6
136				Female	9.1
137	Basheer SM [85]	2019	BRN	Female	5.0
138	Kesu Belani L [80]	2019	MAL	Female	2.0
139	Azizalrahman A [87]	2020	KSA	Female	3.0
140	McGrath RB [88]	2020	IRL	Male	12
141	Shafemaleiq MA [89]	2020	MAL		2.0
142	Sutcu M [90]	2020	TR	Female	2.0
143				Male	0.5

Supplementary table 2:

Characteristics of 36 female subjects affected by genital hair-thread strangulation syndrome. The country of publication is identified using the International vehicle registration code.

Case	First author	Year of publication	Country	Age (years)
1	Chapman HL [91]	1968	CDN	9
2	Summers JL [93]	1973	USA	11
3	Webley JA [94]	1973	USA	7
4	Press S [92]	1980	USA	8
5	Beasley S.W [22]	1991	AUS	8
6	Rich MA [95]	1999	USA	5
7	Sylvestzrak MS [96]	2000	USA	5
8	Kuo JH [97]	2001	USA	14
9	Golshevsky J [100]	2004	AUS	14
10	Klusmann A [36]	2004	D	12
11	Bacon JL [99]	2005	USA	13
12	Osterhoudt KC [101]	2005	USA	8
13	Alverson B [102]	2007	USA	9
14	Serour F [98]	2007	IL	4
15	Cutroni M [49]	2009	I	0.5
16	Pomeranz M [103]	2009	IL	11
17	Stock C [104]	2012	USA	12
18	Bannier MA [105]	2013	NL	6
19	Dua A [106]	2013	USA	12
20	Lee JH [107]	2013	UK	11
22	Ozcakir E [108]	2014	TR	9
22	Panagidis A [109]	2015	GR	14
23				10
24	Parlak M [110]	2015	TR	6
25	Zimmerman LN [111]	2015	USA	5
26	Zhong N [112]	2017	USA	10
27	Ihara T [86]	2018	J	14
28	Roman E [113]	2018	AUS	12
29	Erikci VS [114]	2019	TR	8
30				7
31				9
32	Imam AS [115]	2019	USA	6
33	Diaz-Morales O [116]	2020	E	10
34	Minodier P [117]	2020	F	5
35	Chawla RK [119]	2021	USA	10
36	Adjei N [118]	2021	USA	11

Supplementary table 3:

Characteristics of 36 male subjects affected by genital hair-thread strangulation syndrome. The country of publication is identified using the International vehicle registration code.

Case	First author	Year of publication	Country	Age (months)
1	DeWeese JA [120]	1952	USA	22
2	Alpert JJ [7]	1965	USA	13
3	Farah R [121]	1973	USA	132
4	Kerry RL [12]	1973	USA	2
5				4
6				10
7	Haddad FS [122]	1982	USA	24
8				24
9	McClure WJ [123]	1985	USA	60
10	Sheinfeld J [124]	1985	USA	84
11	Mariani PJ [125]	1986	USA	60
12	Vazquez Rueda F [24]	1996	E	24
13				48
14	Pantuck AJ [126]	1997	USA	36
15	Bangroo AK [127]	2005	IND	120
16	Chaware SM [128]	2006	IND	84
17	Dar NR [129]	2007	PK	60
18	Hussain HM [130]	2008	AUS	72
19	Okeke LI [131]	2008	NGR	108
20	Basner A [132]	2012	D	2
21	Pahwa HS [133]	2013	IND	96
22	Jesus LE [134]	2014	BR	120
23	Dikshit VK [135]	2015	IND	120
24	Gazali ZA [136]	2015	IND	72
25	Zengin K [137]	2015	TR	96
26	Sekhon V [138]	2017	IND	84
27	Sheikh Andalibi MS [139]	2017	IR	54
28	Arshadi H [141]	2019	IR	108
29	Degheili JA [142]	2019	RL	48
30	Ozciit B [140]	2019	TR	48
31	Bouassida K [143]	2020	TN	108
32	Rawls WF [144]	2020	PK	24
33				17
34	Loloi J [147]	2021	USA	96
35	Patterson JW [145]	2022	USA	0.5
36	Rampersad B [146]	2022	TT	24

Supplementary table 4:

Characteristics of 11 individuals affected by non-digital and non-genital hair-thread strangulation syndrome. The country of publication is identified using the International vehicle registration code.

Case	First author	Year of publication	Country	Sex	Age (months)
1	McNeal RM [148]	1987	USA	Male	13
2	Krishna S [149]	2003	USA		3
3	Chegwidden HJ [150]	2005	USA	Female	11
4	Schneider K [151]	2013	USA	Female	84
5	Flores JR [152]	2014	MEX		24
6					48
7	Alghamdi F [153]	2018	KSA	Female	48
8	Wong K [154]	2019	USA	Male	13
9	Chan LYF [156]	2020	HK	Male	21
10	Stansfield J [155]	2020	UK		9
11	Leighton J [157]	2021	UK	Female	36