A rare cause of Sister Mary Joseph's nodule

Rayid Abdulqawi^a, Saqib Ahmad^a, Khaled Ashawesh^b

- ^a Department of Medicine, Princess Royal Hospital, Apley Castle, Telford, UK
- b Diabetes and Endocrinology Department, Walsgrave Hospital, University Hospitals Coventry and Warwickshire NHS Trust, Coventry, UK

Summary

Sister Mary Joseph's nodule is an important physical sign. It usually represents umbilical metastasis of intra abdominal malignancy. Its association with intra abdominal desmoplastic small round cell tumour is extremely rare. We describe a rare case of sister Mary Joseph's nodule associated with desmoplastic small round cell tumour. We

conclude that differential diagnosis of Sister Mary Joseph's nodule should include desmoplastic small round cell tumour, especially in adolescents and young adults.

Key words: Sister Mary Joseph's nodule; desmoplastic small round cell tumour

Case report

A 28-year-old Caucasian male presented with a two-week history of episodic colicky central abdominal pain. The past medical history was significant for ulcerative colitis. On physical examination the abdomen was mildly distended but no mass was palpable. A violaceous periumbilical nodule (figure 1) was consistent with Mary Joseph's nodule, and bilateral firm, hard inguinal lymph nodes were present.

Abdominal and pelvic CT revealed a 5.5 cm lobulated mass lying between the anterior aspect of the rectum and bladder (figure 2, arrow), with several peritoneal deposits but no ascites. It also showed bilateral obstruc-

tion of both ureters distally with bilateral hydronephrosis.

Histopathological evaluation of inguinal lymph mour with small cell morphology. The tissue diagnosis of desmoplastic small round cell tumor (DSRCT) was confirmed on the basis of a characteristic t(11;12)(p13;q12) cytogenetic translocation.

The patient underwent bilateral nephrostomy. The neoplasm was inoperable and he is currently being treated by systemic chemotherapy.



Figure 1
Sister Mary Joseph's nodule.



Figure 2

Axial CT of abdomen and pelvis shows the mass in the rectovesical pouch (white arrow).

Discussion

Sister Mary Joseph's nodule is a rare but important clinical sign. It usually represents an umbilical metastasis of an intraabdominal malignancy, with most of the lesions arising from the stomach, large bowel, pancreas and ovary. It is named after Sister Mary Joseph (1856–1939),

Grants or financial Support: none

the superintendent of St Mary's Hospital in Rochester, who noted the association between umbilical nodules observed during skin preparation for surgery and metastatic intraabdominal cancer confirmed at operation [1]. Its association with intraabdominal DSRCT has rarely been described. To the best of our knowledge there have been only two previous reports in the literature of Sister Mary Joseph's nodule as a presenting sign of DSRCT [2, 3].

DSRCT is a rare, highly aggressive cancer usually affecting male adolescents and young adults. It was first described in 1989 by Gerlad and Rosai [4].

It is usually intraabdominal, often with diffuse peritoneal implants. Common presenting symptoms and signs are abdominal pain, palpable abdominal mass, abdominal distension, vomiting, hepatomegaly and palpable lymph nodes [5, 6]. Other associated features are intestinal and ureteral obstruction [6].

Histologically the tumour is characterised by well-defined nests of small round blue cells separated by abundant desmoplastic stroma. Cytogenetic study reveals a characteristic reciprocal chromosomal translocation, t(11;12)(p13;q12) [7].

CT findings include bulky intraabdominal soft-tissue masses involving omental and serosal surfaces, without a distinct organ of origin; solid, dominant, heterogeneous pelvic masses in the retrovesical or rectouterine spaces; and concurrent metastases common at the time of diagnosis, particularly those involving lymph nodes and liver [8].

Regardless of the treatment modality (chemotherapy, surgery or radiotherapy), the disease is almost uniformly fatal.

DSRCT should be considered in the differential diagnosis of Sister Mary Joseph's nodule, especially in a young male patient with an intraabdominal mass. Diagnosis is based on recognition of characteristic CT, histopathological and cytogenetic findings.

Correspondence:
Dr. Rayid Abdulqawi
Department of Medicine
Princess Royal Hospital
Apley Castle
Telford TF1 6TF
UK
E-Mail: rayid_abdulqawi@hotmail.com

References

- 1 Hill M, O'Leary JP. Vignettes in medical history. Sister Mary Joseph and her node. Am Surg. 1996;62(4):328–9.
- 2 Albano EA, Kanter J. Images in Clinical Medicine. Sister Mary Joseph's Nodule. N Engl J Med. 2005;352(18):1913.
- 3 Doros L, Kaste SC, Rodriguez-Galindo. Sister Mary Joseph's Nodule as Presenting Sign of a Desmoplastic Small Round Cell Tumor. Pediatr Blood Cancer. 2006; Epub ahead of print.
- 4 Gerald WL, Rosai J. Desmoplastic small round cell tumor with divergent differentiation. Pediatr Pathol. 1989;9:177–83.
- 5 Livaditi E, Mavridis G, Soutis M, et al. Diffuse intraabdominal desmoplastic small round cell tumor: a ten year experience. Eur J Pediatr Surg. 2006;16(6):423–7.
- 6 Hassan I, Shyyan R, Donohue JH, et al. Intraabdominal desmoplastic small round cell tumors: a diagnostic and therapeutic challenge. Cancer. 2005;104(6):1264–70.
- 7 Sawyer JR, Tryka AF, Lewis JM. A novel reciprocal chromosome translocation t(11;12)(p13;q12) in an intraabdominal desmoplastic small round cell tumor. Am J Surg Pathol. 1992;16:411–6.
- 8 Bellah R, Suzuki-Bordalo L, Brecher E, et al. Desmoplastic small round cell tumor in the abdomen and pelvis: report of CT findings in affected children and young adults. AJR Am J Roentgenol. 2005;184(6):1910–4.

Established in 1871

Formerly: Schweizerische Medizinische Wochenschrift

Swiss Medical Weekly

Official journal of the Swiss Society of Infectious diseases, the Swiss Society of Internal Medicine and the Swiss Respiratory Society

The many reasons why you should choose SMW to publish your research

What Swiss Medical Weekly has to offer:

- SMW's impact factor has been steadily rising. The 2005 impact factor is 1.226.
- Open access to the publication via the Internet, therefore wide audience and impact
- Rapid listing in Medline
- LinkOut-button from PubMed with link to the full text website http://www.smw.ch (direct link from each SMW record in PubMed)
- No-nonsense submission you submit a single copy of your manuscript by e-mail attachment
- Peer review based on a broad spectrum of international academic referees
- Assistance of our professional statistician for every article with statistical analyses
- Fast peer review, by e-mail exchange with the referees
- Prompt decisions based on weekly conferences of the Editorial Board
- Prompt notification on the status of your manuscript by e-mail
- Professional English copy editing
- No page charges and attractive colour offprints at no extra cost

Editorial Board

Prof. Jean-Michel Dayer, Geneva

Prof. Peter Gehr, Berne

Prof. André P. Perruchoud, Basel

Prof. Andreas Schaffner, Zurich

(Editor in chief)

Prof. Werner Straub, Berne

Prof. Ludwig von Segesser, Lausanne

International Advisory Committee

Prof. K. E. Juhani Airaksinen, Turku, Finland Prof. Anthony Bayes de Luna, Barcelona, Spain

Prof. Hubert E. Blum, Freiburg, Germany

Prof. Walter E. Haefeli, Heidelberg, Germany

Prof. Nino Kuenzli, Los Angeles, USA

Prof. René Lutter, Amsterdam,

The Netherlands

Prof. Claude Martin, Marseille, France

Prof. Josef Patsch, Innsbruck, Austria

Prof. Luigi Tavazzi, Pavia, Italy

We evaluate manuscripts of broad clinical interest from all specialities, including experimental medicine and clinical investigation.

We look forward to receiving your paper!

Guidelines for authors: http://www.smw.ch/set_authors.html



All manuscripts should be sent in electronic form, to:

EMH Swiss Medical Publishers Ltd. SMW Editorial Secretariat Farnsburgerstrasse 8 CH-4132 Muttenz

Manuscripts: su Letters to the editor: le Editorial Board: re Internet: h

submission@smw.ch letters@smw.ch red@smw.ch http://www.smw.ch