

Remodelling in young sheep: a histological study of experimentally produced defects of the TMJ

Orhan Güven^a, Murat Metin^b, Ahmet Keskin^a

^a Department of Oral and Maxillofacial Surgery, School of Dentistry, University of Ankara, Ankara, Turkey

^b Department of Oral and Maxillofacial Surgery, School of Dentistry, University of Ondokuz Mayıs, Samsun, Turkey

Summary

Questions under study: In this study, experimentally induced surgical defects of the mandibular condyle were evaluated histologically 3, 6 and 9 months postoperatively to investigate the capacity for remodelling which is one of the most important features of the temporomandibular joint.

Methods: In three healthy sheep standardised surgical defects were produced on each right condylar head. The left temporomandibular joints served as controls. No treatment was performed on the surgical defects and function was not restricted postoperatively. At the end of the experi-

mental period the sheep were sacrificed and a histological evaluation was carried out.

Results: Histologically, no disturbance was observed during the healing process in bone and cartilage. Although function was not limited throughout the study, it was observed that the experimental surgical defects had healed completely.

Conclusions: This result points to the condyle's high capacity for repair and remodelling.

Key words: mandibular condyle; remodelling; sheep

Introduction

Temporomandibular joint dysfunction may be caused by an abnormal relationship between disc and condyle. The resolution of this condition is often achieved by remodelling and adaptation rather than actual correction of the anomaly [1]. Trauma to the chin may cause injury to the articular tissues, resulting in condylar fractures and posttraumatic arthritis [2]. Condylar fractures are often followed by displacement of the condylar head. During the healing period, remodelling of the joint may occur. In children, extensive remodelling resulting in almost complete restoration of

joint anatomy takes place in many cases [3]. Remodelling has also been observed in adult patients, but to a lesser degree [4]. Trauma to the TMJ, including microtrauma, has been suggested as a possible cause of osteoarthrotic changes of the condylar head. In this context there is still debate as to whether the degenerative changes to the TMJ occur as a result of damage to the mandibular condyle cartilage or of damage to the articular disc.

The objective of the present study is to analyse remodelling of the TMJ after surgical trauma in an experimental model.

Methods

Three young adult sheep aged 12 months and weighing approximately 50 kg were used. The right-side joints were surgically altered and the left-side joints served as controls. Anaesthesia (4 mg ketamine HCL and 0.05 mg xylazine HCL per kilogram) was induced intravenously into the external jugular vein. The skin over the TMJ region was shaved and prepared with betadine solution. Fol-

lowing a preauricular incision, the condylar process was exposed and the inferior joint space opened by a horizontal incision through the capsule. A 0.5 cm wide and 0.5 cm deep mediolateral groove was cut at the top of the condyle midline of the articular surface (Fig. 1) with a 1.5 mm diameter fissure bur using copious amounts of saline for cooling. The condylar surfaces were irrigated with saline

and the joint capsule and overlying tissues repaired in layers. The animals' preoperative and postoperative diet was the same. Postoperatively 2 ml of clindamycin was given twice daily for 7 days intramuscularly. No wound became infected and all the sheep did well following surgery.

The sheep were killed by anaesthetic overdose after 3 months, 6 months and 9 months. The specimens were placed in 10% neutral formaldehyde solution and decalcified in nitric acid solution. All specimens were embedded in paraffin. Serial sections were cut at 5 μ m and stained with haematoxylin and eosin.

Results

After creation of the surgical trauma the animals continued to gain weight. The histopathological findings in all TMJs were as follows:

1. At the third postoperative month, microscopic examination of the joint surface revealed that the defect extended to the cancellous bone and was filled with connective tissue containing blood vessels. Since the defect was distant from the attachment of the synovial membrane, it was thought that the connective tissue cells originated from the osteogenic cells surrounding the bony trabeculae. A surplus of connective tissue mass developed at the defect and caused doming at the joint surface (Fig. 2).
2. At the sixth postoperative month the defect was filled with connective tissue. At some sites

formation of cartilage tissue from the base of the defect was observed (Fig. 3).

3. At the ninth postoperative month histopathological examination showed persistent slight doming at the surface of the defect site. The greater part of the defect was filled with cartilage. Only the central part of a small area at the joint surface consisted of loose connective tissue. Undifferentiated cells resembling cartilage cells were seen parallel to the joint surface, and cellularity was increased close to the surface. Spongy bone developed via endochondral ossification at the base of the defect (Fig. 4).

Figure 1

Surgically created defect on the condylar head.



Figure 2

3 months: Infiltration of the prepared defect and surrounding joint surface with connective tissue containing blood vessels. H E. Magnification $\times 80$.




Figure 3

6 months: Formation of cartilage tissue from the base of the defect to the joint surface. H E. Magnification $\times 80$.





Figure 4

9 months: Appearance of the defect of joint surface. A major portion of the defect was filled with cartilage tissue. The excessive portion at joint surface consisted of loose connective tissue. However, especially in the upper portions, differentiation to cartilage tissue seemed more likely. H E. Magnification $\times 80$.



Discussion

The condylar portion of the TMJ is most commonly affected by disease, such as osteoarthritis, trauma or errors of development.

In various clinical and experimental studies it has been found that mandibular condylar cartilage has the capacity to remodel after acute or chronic trauma [2-8]. However, complications subsequent to condylar trauma such as ankylosis, osteoarthritis and mandibular deformities have also been described [9-14].

Although remodelling of the condylar process following condylar fracture has been demonstrated [15-17], the repair mechanisms within the TMJ are poorly understood.

In their study Teixeira et al. [18] showed that displaced condylar process fractures in rats healed by callus formation with concurrent repositioning of the condyle. In man remodelling of the joint occurs during the healing of condylar fractures [15, 19, 20]. In growing patients extensive remodelling, resulting in almost complete restitution of the joint, has been noted in many cases. Remodelling has also been observed in adult patients, but to a lesser degree [4].

Hochman and Laskin [21], using rabbits, reported that surgical defects of the mandibular condyle were completely repaired by both adjacent cartilage and reparative tissue from the marrow space. The difference in the repair process can be ascribed to the animals' differing characteristics: rabbits have greater potential for intrinsic repair of bone and cartilage defects, while humans have less. In this respect sheep are similar to man, and thus the choice of animal is important in these experiments if we wish to compare the biological changes with human disease [22]. Bifano et al. [23] report that goats and sheep are well suited as TMJ models due to their relative lack of protection against anteroposterior forces, their relative lack of posterior joint function and the translatory capacity of the joint. They also have the advantage of being plentiful, inexpensive, easily maintained and easily handled. It was for these reasons that we chose sheep for this study.

In the present study it was found that a surgically created defect of the condyle was replaced 3

months postoperatively by fibro-osseous connective tissue associated with new blood vessel formation. Histologically no alteration was observable in the control joints. However, all three articular discs had thinned. Six months postoperatively chondrocyte proliferation and a decrease in the quantity of connective tissue were noted. The chondrocytes probably derived from marrow. In similar fashion to our results, the defect created in the marmoset by Robinson [7] was filled with an increasingly cellular blastema forming new collagen fibres. He reported that these cells appeared to be in continuity with the subchondral bone. According to Ishimaru et al. [24] the residual articular cartilage has less potential for wound repair, with greater potential existing in the bone marrow.

It has been suggested that 90 days is too short a period for the advanced stages to develop, nor do major changes take place over 6 months [22]. However, our studies showed differences between 6 and 9 months postoperatively.

The study's purpose was to observe the remodelling capability of condylar cartilage in the repair of standardised full-thickness articular bone and cartilage defects. It illustrates the effect of a situation similar to an intraarticular fracture of the condyle, either a microfracture from repeated overloading or a fracture from acute mandibular trauma.

Although the functioning of the TMJ was not limited throughout this study, complete healing of the experimental surgical defects was observed. This points to the condyle's high capacity for repair and remodelling, a factor which should be considered in the management of trauma and other disorders of the TMJ.

Correspondence:

Dr. Orhan Güven

Yeşilyurt sok 24/15

A. Ayrancı

06690 Ankara

Turkey

E-Mail: oguvenc@dentistry.ankara.edu.tr

References

- 1 Blaustein DI, Scapino RP. Remodelling of the temporomandibular joint disk and posterior attachment in disk displacement specimens in relation to glycosaminoglycan content. *Plast Reconstr Surg* 1986;78:756-4.
- 2 Luz JGC, Jaeger RG, De Araujo VC, De Rezende RJV. The effect of immobilisation on the rabbit temporomandibular joint. *Int J Oral Maxillofac Surg* 1991;20:48-2.
- 3 Güven O, Keskin A. Remodelling following condylar fractures in children. *J Cranio-Maxillofac Surg* 2001;29:232-7.
- 4 Mongini F. Remodelling of the mandibular condyle in the adult and relationship to the condition of the dental arches. *Acta Anat* 1972;82:437-3.
- 5 Avrahami E, Frishman E, Weiss-Peretz J, Horowitz I. Computed tomography of healing condylar fractures with some clinical correlations. *Clin Radiol* 1993;47:269-3.
- 6 Leake D, Doykos J, Habal MB, Murray JE. Long term follow-up of fractures of the mandibular condyle in children. *Plast Reconstr Surgery* 1971;47:127-1.
- 7 Robinson PD. Articular cartilage of the temporomandibular joint: Can it regenerate? *Ann R Coll Surg* 1993;75:231-6.
- 8 Yasuoka T, Oka N. Histomorphometric study of trabecular bone remodelling during condylar process fracture healing in the growing period. *J Oral Maxillofac Surg* 1991;49:981-8.

- 9 Norholt SE, Krishnan V, Sindet-Pedersen S, Jensen AI. Pediatric condylar fractures: A long term follow-up study of 55 patients. *J Oral Maxillofac Surg* 1993;51:1302-9.
- 10 Proffit WR, Vig KWL, Turvey TA. Early fracture of the mandibular condyles: Frequently an unsuspected cause of growth disturbances. *Amer J Orthod* 1980;78:1-4.
- 11 Sawhney CP. Bony ankylosis of the temporomandibular joint: Follow-up of 70 patients treated with arthroplasty and acrylic spacer interposition. *Plast Reconstr Surg* 1986;77:29-8.
- 12 Wu XG, Hong M, Sun KH. Severe osteoarthritis after fracture of the mandibular condyle: A clinical and histological study of seven patients. *J Oral Maxillofac Surg* 1994;52:138-2.
- 13 Güven O. A clinical study on temporomandibular joint ankylosis. *Auris Nasus Larynx* 2000;27:27-3.
- 14 Güven O. Acrylic spacer in treatment of TMJ ankylosis (a technical report). *Balkan J Stomatology* 2000;4:40-1.
- 15 Feifel H, Albert-Deumlich J, Riediger D. Long term follow up of subcondylar fractures in children by electronic computer-assisted recording of condylar movements. *Int J Oral Maxillofac Surg* 1992;21:70-6.
- 16 Sahm G, Schuknecht B, Eberhardt K. Zur Remodellation nach Kondylusluxationsfracturen bei Heranwachsenden. *Dtsch Zahnarzt* 1990;45:403-5.
- 17 Zou ZJ, Wu WT, Sun GX, et al. Remodelling of the temporomandibular joint after conservative treatment of condylar fractures. *Dentomaxillofac Radiol* 1987;16:91-8.
- 18 Teixeira ACB, Luz JGC, Aaujo VC, Araujo NS. Healing of the displaced condylar process fracture: an experimental study. *J of Cranio-maxillofac Surg* 1998;26:326-0.
- 19 Kahl-Nieke B, Fischbach R, Gerlach KL. CT analysis of temporomandibular joint state in children 5 years after functional treatment of condylar fractures. *Int J Oral Maxillofac Surg* 1994;23:332-7.
- 20 Kahl-Nieke B, Fischbach R, Gerlach KL. Temporomandibular joint morphology in children after treatment of condylar fractures with functional appliance therapy: a follow up study using spiral computed tomography. *Dentomaxillofac Radiol* 1995;24:37-5.
- 21 Hochman LS, Laskin DM. Repair of surgical defects in the articular surface of the rabbit mandibular condyle. *Oral Surg* 1965;19:534-2.
- 22 Ishimaru J, Goss AN. A model for osteoarthritis of the temporomandibular joint. *J Oral Maxillofac Surg* 1992;50:1191-5.
- 23 Bifano C, Hubbard G, Ehler W. A comparison of the form and function of the human, monkey, and goat temporomandibular joint. *J Oral Maxillofac Surg* 1994;52:272-5.
- 24 Ishimaru JI, Kurita K, Handa Y, Goss AN. Effect of marrow perforation on the sheep temporomandibular joint. *Int J Oral Maxillofac Surg* 1992;21:239-2.

The many reasons why you should choose SMW to publish your research

What Swiss Medical Weekly has to offer:

- SMW's impact factor has been steadily rising, to the current 1.537
- Open access to the publication via the Internet, therefore wide audience and impact
- Rapid listing in Medline
- LinkOut-button from PubMed with link to the full text website <http://www.smw.ch> (direct link from each SMW record in PubMed)
- No-nonsense submission – you submit a single copy of your manuscript by e-mail attachment
- Peer review based on a broad spectrum of international academic referees
- Assistance of our professional statistician for every article with statistical analyses
- Fast peer review, by e-mail exchange with the referees
- Prompt decisions based on weekly conferences of the Editorial Board
- Prompt notification on the status of your manuscript by e-mail
- Professional English copy editing
- No page charges and attractive colour offprints at no extra cost

Editorial Board

Prof. Jean-Michel Dayer, Geneva
 Prof. Peter Gehr, Berne
 Prof. André P. Perruchoud, Basel
 Prof. Andreas Schaffner, Zurich
 (Editor in chief)
 Prof. Werner Straub, Berne
 Prof. Ludwig von Segesser, Lausanne

International Advisory Committee

Prof. K. E. Juhani Airaksinen, Turku, Finland
 Prof. Anthony Bayes de Luna, Barcelona, Spain
 Prof. Hubert E. Blum, Freiburg, Germany
 Prof. Walter E. Haefeli, Heidelberg, Germany
 Prof. Nino Kuenzli, Los Angeles, USA
 Prof. René Lutter, Amsterdam,
 The Netherlands
 Prof. Claude Martin, Marseille, France
 Prof. Josef Patsch, Innsbruck, Austria
 Prof. Luigi Tavazzi, Pavia, Italy

We evaluate manuscripts of broad clinical interest from all specialities, including experimental medicine and clinical investigation.

We look forward to receiving your paper!

Guidelines for authors:

http://www.smw.ch/set_authors.html

Impact factor Swiss Medical Weekly



All manuscripts should be sent in electronic form, to:

EMH Swiss Medical Publishers Ltd.
 SMW Editorial Secretariat
 Farnsburgerstrasse 8
 CH-4132 Muttenz

Manuscripts: submission@smw.ch
 Letters to the editor: letters@smw.ch
 Editorial Board: red@smw.ch
 Internet: <http://www.smw.ch>